

Movements and spatial use of odontocetes in the western main Hawaiian Islands: results from satellite-tagging and photo-identification off Kaua‘i and Ni‘ihau in July/August 2011

Robin W. Baird¹, Daniel L. Webster¹, Gregory S. Schorr¹,
Jessica M. Aschettino¹, Antoinette M. Gorgone², Sabre D. Mahaffy¹

¹Cascadia Research Collective

218 ½ W. 4th Avenue

Olympia, WA 98501

²NOAA Southeast Fisheries Science Center

101 Pivers Island Road

Beaufort, NC 28516

Annual progress report under Grant No. N00244-10-1-0048

from the Naval Postgraduate School

June 7, 2012

Summary

Although considerable information is available on residency patterns and spatial use of odontocetes in the eastern half of the Hawai'i Range Complex (HRC), much less is known about odontocetes in the western half of the HRC. In the second year of a three-year effort in the western main Hawaiian Islands we undertook surveys off Kaua'i and Ni'ihau in July/August 2011, to examine spatial use and residency patterns using satellite tags, to provide visual verification of acoustically-detected odontocetes on the Pacific Missile Range Facility (PMRF), and to obtain individual identification photographs and biopsy samples for assessment of population identity and structure. During 18 days of field effort we covered 1,972 km of trackline and had 65 encounters with five species of odontocetes. Twenty-four of the encounters, of three species, were cued by acoustic detections from the Marine Mammal Monitoring on Navy Ranges (M3R) system, thus providing species verifications for future use of the M3R system on the PMRF range. During the 65 encounters we obtained 22,645 photos for individual and species identification, and collected 48 biopsy samples for genetic analyses. One encounter with a group of four killer whales was only the second encounter with this species in 12 years of directed field surveys in Hawaiian waters. Photos from that encounter were compared to our photo-identification catalog but no matches were found, further suggesting that there is no population of this species resident to the Hawaiian Islands. There were three encounters with a lone pantropical spotted dolphin, each time in association with a group of spinner dolphins. Photos of this individual matched to a spotted dolphin identified off Kaua'i in 2004 and in 2005, both times with spinner dolphins, suggesting this individual may be part of a long-term association with spinner dolphins. Four satellite tags were deployed; three on rough-toothed dolphins and one on a bottlenose dolphin. These are the first tag deployments on either species in Hawaiian waters and the first deployments of satellite tags on free-ranging rough-toothed dolphins anywhere in the world. Rough-toothed dolphin tag data were obtained over periods from 7.6 to 18.5 days. Over these periods the three rough-toothed dolphins moved cumulative horizontal distances ranging from 573 to 1,295 km, yet remained an average distance from the tagging locations of from 10.4 to 13.9 km. Median depths used by the three rough-toothed dolphins ranged from 816 to 1,107 m, with median distance from shore ranging from 11.6 to 12.2 km. Two of the three individuals had been previously photo-identified off Kaua'i (in 2007 or 2008), and all link by association with the resident population from Kaua'i and Ni'ihau.

Movement and habitat use data were obtained over a 34-day period for the satellite-tagged bottlenose dolphin. During this time the individual remained associated with the island of Kaua'i using waters with a median depth of 82 m. Although this individual had not been previously photo-identified, others from the group it was in had been previously documented off Kaua'i and/or Ni'ihau in 2003-2005, suggesting it is part of the island-resident population. Overall these efforts provide the first unbiased movement and habitat use data for both species in Hawaiian waters.

Introduction

Considerable information is available on residency patterns and spatial use of a number of species of odontocetes in the eastern half of the Hawai'i Range Complex (HRC), particularly off the island of Hawai'i (e.g., McSweeney et al. 2007, 2009; Baird et al. 2008a, 2008b, 2009, 2010, 2011; Schorr et al. 2009; Aschettino et al. 2011). Favorable working conditions have resulted in a concentration of research activities off the island, where the presence of very deep water (>2,000 m) close to shore has facilitated research with a number of typically deep-water species, as well as with shallow-water species. One of the main findings of this work is that there are resident populations of more than half of the species of odontocetes found off the island of Hawai'i, including short-finned pilot whales, pygmy killer whales, melon-headed whales, common bottlenose dolphins, rough-toothed dolphins, Blainville's beaked whales, Cuvier's beaked whales, and dwarf sperm whales (e.g., McSweeney et al. 2007, 2009; Baird et al. 2008a, 2009, 2010, 2011; Schorr et al. 2009; Aschettino et al. 2011; Courbis 2011; Mahaffy 2012; Cascadia Research Collective, unpublished data). In addition to these resident populations, individuals of some species are known to regularly move among the islands (e.g., false killer whales; Baird et al. 2008b, 2010), and at least one has two populations that use the area (melon-headed whales, which have both a resident population to the island of Hawai'i and a population that moves among all of the main Hawaiian Islands and into offshore waters; Aschettino et al. 2011, Woodworth et al. 2011).

Less is known about residency and spatial use of odontocetes in the western half of the HRC. The smaller size of the islands (Kaua'i, Ni'ihau, O'ahu) result in smaller lee areas and thus less ideal working conditions, and the shallower slopes to the islands mean that deep water is

further offshore and thus in less protected areas. Most of the research that has been done has focused on photo-identification and biopsy sampling (e.g., Baird et al. 2003, 2006a, but see Baird et al. 2008c, 2011), and has identified the existence of some island-specific populations of more commonly-encountered species, like rough-toothed dolphins off Kaua'i/Ni'ihau (Baird et al. 2008a) and common bottlenose dolphins (hereafter bottlenose dolphins) off both Kaua'i/Ni'ihau and O'ahu (Baird et al. 2009; Martien et al. 2011). However, there is a paucity of information available for less-frequently encountered and/or deeper-water species. Navy exercises are concentrated in the western half of the HRC, thus there is a need for additional information on residency patterns and spatial use of protected species in that area, to be able to assess the likelihood of exposure to exercises, interpret potential reactions, and assess overall impacts. One of the primary purposes of this project is to address this information gap. A number of tools are used to accomplish this, in particular deployment of medium-term Low Impact Minimally Percutaneous External-electronics Transmitter (LIMPET) satellite tags on a number of species of odontocetes. Importantly, photo-identification results are also used to interpret spatial use patterns, building on long-term photo-identification catalogs compiled from throughout the main Hawaiian Islands, and collection of genetic samples to contribute to studies of population structure. In the first year of effort under this grant field activities were undertaken off O'ahu (in October 2010) and Kaua'i (in February 2011), with considerable new information obtained on movements and habitat use of a number of species of odontocetes (Baird et al. 2011). This report summarizes results from the second year of this effort, with field activities undertaken in July and August 2011 off the islands of Kaua'i and Ni'ihau. In addition to support from N45 and the Naval Postgraduate School, these field efforts were supported in part by funding from Commander, Pacific Fleet.

Methods

Surveys were undertaken off the island of Kaua'i for 18 days between 21 July and 8 August 2011. Several research vessels were used: a 27' Boston Whaler outfitted with a tower and bow pulpit; a 24' rigid-hulled Zodiac; and a 32' Sea Cat. When the PMRF range was open to vessel traffic, survey efforts were coordinated with the Marine Mammal Monitoring on Navy Ranges (M3R) program with the research vessel directed to areas with acoustic identifications of odontocetes. When access to the range was closed surveys were undertaken to the south, west or

east of the range. Observers scanned 360 degrees around the research vessel, which transited typically at speeds of 15-30 km h⁻¹. Efforts were made to obtain photographs of all individuals in groups of odontocetes encountered and biopsy samples of most species. A laser photogrammetry system using two green dot lasers mounted 15 cm apart (Webster et al. 2010) was used to obtain dorsal fin measurements when possible, to assist in assessing age class of individuals.

Satellite tagging was to be undertaken with any of a number of priority target species encountered, including beaked whales, sperm whales, melon-headed whales, false killer whales, short-finned pilot whales, or pygmy killer whales. Depending on encounter rates (and tags deployed) with those species, satellite tags were also to be deployed on lower-priority species such as rough-toothed dolphins and bottlenose dolphins. The satellite tags used were location-only SPOT5 tags (Wildlife Computers, Inc., Redmond, WA), in the LIMPET configuration (Andrews et al. 2009; Schorr et al. 2009; Baird et al. 2010). Attachment darts penetrated 4.2 cm into the dorsal fin. Tags were programmed to transmit for variable periods during the day corresponding to the periods with the best satellite overpasses. Tagged individuals were biopsy sampled when possible, to allow for genetic sexing of individuals (analyses undertaken by R. Albertson, Oregon State University).

Data obtained from the Argos system was processed with the Douglas Argos-Filter v. 7.08 (available at Alaska.usgs.gov/science/biology/spatial/douglas.html) using two independent methods: distance between consecutive locations, and rate and bearings among consecutive movement vectors. Each location is assigned a "location class" by Argos, which reflects the estimated precision of the location, with the most precise locations being classes 3 and 2 (estimated error of <250 m and < 500 m, respectively). We set the Douglas Argos-Filter to automatically retain location classes 3 and 2. Maximum rate of movement was set at 20 km h⁻¹. Depth and distance from shore for all locations which passed the Douglas Argos-filter were determined in ArcGIS v. 9.2 (ESRI, Redlands, California) using a 50 m x 50 m multibeam synthesis bathymetry model from the Hawai'i Mapping Research Group (available at www.soest.hawaii.edu/HMRG/multibeam/index.php).

To determine whether individuals with overlapping tag data were acting in concert or independently, we calculated the straight-line distance (i.e., not taking into account potentially intervening land masses) between pairs of individuals when locations were obtained during a single satellite overpass. We used both the average distances between pairs of individuals and the maximum distance between pairs to assess whether individuals were acting independently.

Photographs of tagged and companion individuals were added and compared to individual photo-identification catalogs for each species maintained by Cascadia Research Collective (e.g., Baird et al. 2006b; Baird et al. 2008a; Baird et al. 2009). Previous sighting histories of individuals within groups were examined to assess whether individuals were: 1) part of resident populations from the areas they were tagged (or whether the group they were in included known resident individuals); 2) potentially part of offshore populations; or 3) individuals moving from other islands. For rough-toothed dolphins, a social network was generated using good or excellent quality photos of all distinctive and very distinctive individuals photo-identified off Kaua'i from 2003 through 2005 (250 identifications, from Baird et al. 2008a) as well as an additional 114 identifications from 2007 through January 2012. Photographs of spinner dolphins were contributed to a catalog held at the NOAA Pacific Islands Fisheries Science Center.

Results and Discussion

In 18 days of field efforts off Kaua'i in July/August we covered 1,972 kilometers of survey trackline with 118.8 hours of survey effort (Figure 1). During this period we encountered 65 different groups of five different species of odontocetes and obtained 22,645 photographs for individual and species identification (Table 1).

The most unusual sighting of the trip was a group of four killer whales, only the second sighting of killer whales in 12 years of field work in the main Hawaiian Islands (see Baird et al. 2006b). Identification photographs of the four killer whales were compared to Cascadia Research Collective's photo-identification catalog for this species in Hawai'i, but no matches were found. Pantropical spotted dolphins are only infrequently seen off Kaua'i (Baird et al. 2003, 2006a; Courbis 2011), and while there were three sightings of pantropical spotted dolphins during this

field effort, all three were of the same individual, seen on three separate days and each time traveling with a group of spinner dolphins in shallow near-shore waters (Figure 1). Photographs of the spotted dolphin were compared with a catalog of this species and this individual matched to a sighting from Kaua'i from February 2004 (M. Deakos, unpublished) and to a sighting from Kaua'i in November 2005 (see Baird et al. 2006a). In both previous cases the spotted dolphin was also with a group of spinner dolphins and was the only spotted dolphin seen, suggesting that this individual may be a semi-permanent associate with spinner dolphins off Kaua'i.

Of the 65 total encounters, 24 were detected based on the M3R acoustic range, with the research vessel directed to the area of the acoustic detection. Of these 24, there were single sightings of spinner dolphins, bottlenose dolphins, and an unidentified odontocete, as well as 21 sightings of rough-toothed dolphins, a species known to be commonly encountered in the study area (Baird et al. 2008a). We collected 48 genetic samples from four of the five species for studies of population structure, including 23 samples from rough-toothed dolphins; these were sent to Oregon State University for a study of rough-toothed dolphin population structure (see Albertson et al. 2011). Sub-samples of all species were sent to the Southwest Fisheries Science Center for curation and analysis.

Four satellite tags were deployed; three on rough-toothed dolphins (Figure 2) and one on a bottlenose dolphin (Table 2). The satellite tag deployments on three rough-toothed dolphins were the first remote deployments of satellite tags on this species anywhere in the world. With attachment durations of up to 18 days these tags provided the first unbiased movement and habitat use data available for this species in Hawaiian waters. Although the minimum cumulative horizontal distance traveled for the three individuals ranged from 573 to 1,295 km, on average the three individuals remained within 10-14 km of their tagging location (Table 3), and repeatedly returned to the general area near where they were tagged (Figures 3, 4). All three individuals remained strongly associated with the islands of Kaua'i and Ni'ihau, with movements largely centered on the PMRF range (Figure 3). Two of the individuals were tagged on the same day in the same group, and they remained relatively close together during the period when tag data from the two overlapped (Figure 5). Photos of all three individuals were compared to our photo-identification catalog, and two of the individuals, HISb0860 (SbTag001, tagged 31

July 2011) and HISb0785 (SbTag003, tagged 2 August 2011) had been previously documented off of Kaua'i; HISb0860 in June 2008 and HISb0785 in October and November 2007 and June 2008. Although the third individual, HISb0908 (SbTag002) had not been previously photographically documented, this individual was with others that had been previously documented off Kaua'i (including HISb0785) and was also re-sighted off Kaua'i in January 2012 (CRC unpublished data). A social network showing distinctive and very distinctive rough-toothed dolphins identified off Kaua'i and Ni'ihau showed that all three tagged individuals are linked by association with the main component of the social network off those islands (Figure 6), although the first individual tagged has not been seen in association with the second or third individual. During the period of tag overlap, the first tagged individual appeared to remain separated from the other two tagged individuals (which were spatially associated), suggesting we obtained data from two, rather than three, independent groups of rough-toothed dolphins (Figure 5).

The satellite tagged bottlenose dolphin was the first of this species remotely tagged with a satellite tag in Hawaiian waters. While photos of this individual did not match any known individuals in our catalog, of the 45 photo-identified individuals from this encounter, 15 individuals had been previously documented off of Kaua'i or Ni'ihau in either 2003, 2004 or 2005, suggesting the tagged individual is part of the resident population from those islands (Baird et al. 2009). This individual remained associated with the island of Kaua'i for the 34 days that locations were received from the tag (Figure 7), using waters with an average depth of 82 m and remaining an average of 2.24 km from shore (Table 3).

Although the samples sizes are small, this effort, combined with efforts in 2010 (Baird et al. 2011) and additional field effort planned for 2012, substantially increase what is known about the residency and habitat use of several species of odontocetes around the western main Hawaiian Islands. Such knowledge is important in providing a stronger basis for assessment of repeated exposure of individuals to Navy exercises, in interpreting individual and/or group reactions, and in determining high-use or ecologically-important areas of occupancy of individuals from the island-associated populations.

Acknowledgements

Field work in July/August 2011 was funded by the Naval Postgraduate School (N45) and by Commander, Pacific Fleet, coordinated by NAVFAC Pacific. We thank Tara Leota, Shawn Hunadi and Kevin Anderson for coming through with vessels when our primary research vessel was in the shop; the M3R crew, Nancy Dimarzio, Ashley Dilley, Scott Fisher, Susan Jarvis, Steve Martin, and Jessica Ward for directing us to acoustic detections; Dave Mortetti of M3R for planning and coordination; Renee Albertson, Ali Bayless, Michele Bane, Meredith Fagan, Paul Furumo, Samantha Hamilton, Oscar Johnson, David Kuhn, Allan Ligon, Morgan Richie, Julie Rivers, Kim Rogers, Lauren Soloman, Jean Souza, Robert Uyeyama, Elisa Weise, Jamie Werthwine, Becky Wilcox, Pam Willis, Kate Winters, and Jochen Zaeschmar for assistance in the field; Don Moses for assistance with vessels; Dan McSweeney and the Wild Whale Research Foundation for coordinating one of the research vessels; Chris Bane for contribution of rough-toothed dolphin photos from 2007; Mark Deakos for contribution of spotted dolphin photos from 2004; Renee Albertson for providing genetic confirmation of sex for two rough-toothed dolphins; Sarah Courbis for finding the initial spotted dolphin match; Damon Holzer of the Northwest Fisheries Science Center for GIS analyses; and Michele Bane for assistance with logistics off the water. Research was undertaken under NMFS Scientific Research Permits No. 731-1774, 774-1714 and 15330.

Literature Cited

- Albertson, G.R., M. Oremus, R.W. Baird, K.K. Martien, M.M. Poole, R.L. Brownell Jr., F. Cipriano and C.S. Baker. 2011. Staying close to home: genetic analyses reveal insular population structure for the pelagic dolphin *Steno bredanensis*. Presentation at the 19th Biennial Conference on the Biology of Marine Mammals, Tampa, FL, Nov-Dec 2011.
- Andrews, R.D., R.L. Pitman, and L.T. Balance. 2008. Satellite tracking reveals distinct movement patterns for Type B and Type C killer whales in the southern Ross Sea, Antarctica. *Polar Biology* 31, 1461-1468.
- Aschettino, J.M., R.W. Baird, D.J. McSweeney, D.L. Webster, G.S. Schorr, J.L. Huggins, K.K. Martien, S.D. Mahaffy, and K.L. West. 2011. Population structure of melon-headed whales (*Peponocephala electra*) in the Hawaiian Archipelago: evidence of multiple populations based on photo-identification. *Marine Mammal Science* doi: 10.1111/j.1748-7692.2011.00517.x
- Baird, R.W., D.J. McSweeney, D.L. Webster, A.M. Gorgone and A.D. Ligon. 2003. Studies of odontocete population structure in Hawaiian waters: results of a survey through the main Hawaiian Islands in May and June 2003. Report prepared under Contract No. AB133F-02-CN-0106 from NOAA, 7600 Sand Point Way N.E., Seattle, WA 98115 USA. Available from www.cascadiaresearch.org/hawaii/publications.htm
- Baird, R.W., G.S. Schorr, D.L. Webster, S.D. Mahaffy, A.B. Douglas, A.M. Gorgone, and D.J. McSweeney. 2006a. A survey for odontocete cetaceans off Kaua'i and Ni'ihau, Hawai'i, during October and November 2005: evidence for population structure and site fidelity. Report to Pacific Islands Fisheries Science Center, NOAA Fisheries, under Order No. AB133F05SE5197 with additional support from the Marine Mammal Commission and Dolphin Quest. Available from www.cascadiaresearch.org/hawaii/publications.htm
- Baird, R.W., D.J. McSweeney, C. Bane, J. Barlow, D.R. Salden, L.K. Antoine, R.G. LeDuc, and D.L. Webster. 2006b. Killer whales in Hawaiian waters: information on population identity and feeding habits. *Pacific Science* 60:523-530.

- Baird, R.W., D.L. Webster, S.D. Mahaffy, D.J. McSweeney, G.S. Schorr and A.D. Ligon. 2008a. Site fidelity and association patterns in a deep-water dolphin: rough-toothed dolphins (*Steno bredanensis*) in the Hawaiian Archipelago. *Marine Mammal Science* 24:535-553.
- Baird, R.W., A.M. Gorgone, D.J. McSweeney, D.L. Webster, D.R. Salden, M.H. Deakos, A.D. Ligon, G.S. Schorr, J. Barlow and S.D. Mahaffy. 2008b. False killer whales (*Pseudorca crassidens*) around the main Hawaiian Islands: long-term site fidelity, inter-island movements, and association patterns. *Marine Mammal Science* 24:591-612.
- Baird, R.W., G.S. Schorr, D.L. Webster, D.J. McSweeney, M.B. Hanson, and R.D. Andrews. 2008c. Multi-species cetacean satellite tagging to examine movements in relation to the 2008 Rim-of-the-Pacific (RIMPAC) naval exercise. A quick look report on the results of tagging efforts undertaken under Order No. D1000115 from the Woods Hole Oceanographic Institution. Available from www.cascadiaresearch.org/hawaii/publications.htm
- Baird, R.W., A.M. Gorgone, D.J. McSweeney, A.D. Ligon, M.H. Deakos, D.L. Webster, G.S. Schorr, K.K. Martien, D.R. Salden, and S.D. Mahaffy. 2009. Population structure of island-associated dolphins: evidence from photo-identification of common bottlenose dolphins (*Tursiops truncatus*) in the main Hawaiian Islands. *Marine Mammal Science* 25:251-274.
- Baird, R.W., G.S. Schorr, D.L. Webster, D.J. McSweeney, M.B. Hanson, and R.D. Andrews. 2010. Movements of satellite-tagged false killer whales around the main Hawaiian Islands. *Endangered Species Research* 10:107-121.
- Baird, R.W., G.S. Schorr, D.L. Webster, S.D. Mahaffy, J.M. Aschettino and T. Cullins. 2011. Movements and spatial use of satellite-tagged odontocetes in the western main Hawaiian Islands: results of field work undertaken off O'ahu in October 2010 and Kaua'i in February 2011. Annual progress report under Grant No. N00244-10-1-0048 from the Naval Postgraduate School. Available from www.cascadiaresearch.org/hawaii/publications.htm
- Courbis, S.S. 2011. Population structure of island-associated pantropical spotted dolphins (*Stenella attenuata*) in Hawaiian waters. Ph.D. Dissertation, Portland State University. 173 pp. Available from www.cascadiaresearch.org/hawaii/publications.htm
- Mahaffy, S.D. 2012. Site fidelity, associations and long-term bonds of short-finned pilot whales off the island of Hawai'i. M.Sc. Thesis, Portland State University. 151 pp. Available from www.cascadiaresearch.org/hawaii/publications.htm
- Martien, K.K., R.W. Baird, N.M. Hedrick, A.M. Gorgone, J.L. Thieleking, D.J. McSweeney, K.M. Robertson, and D.L. Webster. 2011. Population structure of island-associated dolphins: evidence from mitochondrial and microsatellite markers for common bottlenose dolphins (*Tursiops truncatus*) around the main Hawaiian Islands. *Marine Mammal Science* doi: 10.1111/j.1748-7692.2011.00506.x
- McSweeney, D.J., R.W. Baird and S.D. Mahaffy. 2007. Site fidelity, associations and movements of Cuvier's (*Ziphius cavirostris*) and Blainville's (*Mesoplodon densirostris*) beaked whales off the island of Hawai'i. *Marine Mammal Science* 23:666-687.
- McSweeney, D.J., R.W. Baird, S.D. Mahaffy, D.L. Webster, and G.S. Schorr. 2009. Site fidelity and association patterns of a rare species: pygmy killer whales (*Feresa attenuata*) in the main Hawaiian Islands. *Marine Mammal Science* 25:557-572.

- Schorr, G.S., R.W. Baird, M.B. Hanson, D.L. Webster, D.J. McSweeney and R.D. Andrews. 2009. Movements of satellite-tagged Blainville's beaked whales off the island of Hawai'i. *Endangered Species Research* 10:203-213.
- Webster, T., S. Dawson and E. Slooten. 2010. A simple laser photogrammetry technique for measuring Hector's dolphins (*Cephalorhynchus hectori*) in the field. *Marine Mammal Science* 26:296-308.
- Woodworth, P.A., G.S. Schorr, R.W. Baird, D.L. Webster, D.J. McSweeney, M.B. Hanson, R.D. Andrews and J.J. Polovina. 2011. Eddies as offshore foraging grounds for melon-headed whales (*Peponocephala electra*). *Marine Mammal Science* doi: 10.1111/j.1748-7692.2011.00509.x.

Table 1. Odontocete sightings off Kaua'i/Ni'ihau in July/August 2011 (continued on next page)

Species	Date	Enc. #	Group size	# photos	# genetic samples	# tag deployments
Bottlenose dolphin	22-Jul-11	1	7	121	1	0
Bottlenose dolphin	22-Jul-11	3	7	414	1	0
Bottlenose dolphin	25-Jul-11	1	1	0	0	0
Bottlenose dolphin	25-Jul-11	2	5	287	2	0
Bottlenose dolphin	25-Jul-11	3	16	317	2	0
Bottlenose dolphin	25-Jul-11	4	15	123	0	0
Bottlenose dolphin	27-Jul-11	4	25	364	0	0
Bottlenose dolphin	28-Jul-11	1	10	652	2	0
Bottlenose dolphin	28-Jul-11	2	200	1253	4	0
Bottlenose dolphin	31-Jul-11	4	4	38	0	0
Bottlenose dolphin	03-Aug-11	1	45	2649	6	1
Bottlenose dolphin	05-Aug-11	2	2	167	1	0
Bottlenose dolphin	06-Aug-11	1	2	57	0	0
Killer whale	24-Jul-11	5	4	779	0	0
Pantropical spotted dolphin	04-Aug-11	2	1	10	1	0
Pantropical spotted dolphin	05-Aug-11	4	1	10	0	0
Pantropical spotted dolphin	08-Aug-11	4	1	10	0	0
Rough-toothed dolphin	21-Jul-11	1	4	53	0	0
Rough-toothed dolphin	21-Jul-11	2	2	146	1	0
Rough-toothed dolphin	21-Jul-11	3	5	219	1	0
Rough-toothed dolphin	21-Jul-11	4	1	0	0	0
Rough-toothed dolphin	21-Jul-11	5	65	1783	2	0
Rough-toothed dolphin	23-Jul-11	1	5	223	1	0
Rough-toothed dolphin	23-Jul-11	2	9	166	1	0
Rough-toothed dolphin	24-Jul-11	1	2	0	0	0
Rough-toothed dolphin	24-Jul-11	2	5	222	0	0
Rough-toothed dolphin	24-Jul-11	3	5	275	2	0
Rough-toothed dolphin	24-Jul-11	4	5	87	0	0
Rough-toothed dolphin	24-Jul-11	6	1	0	0	0
Rough-toothed dolphin	24-Jul-11	7	2	78	1	0
Rough-toothed dolphin	26-Jul-11	1	6	400	1	0
Rough-toothed dolphin	27-Jul-11	1	2	0	0	0
Rough-toothed dolphin	27-Jul-11	2	1	0	0	0
Rough-toothed dolphin	27-Jul-11	3	3	517	1	0
Rough-toothed dolphin	28-Jul-11	4	10	497	0	0
Rough-toothed dolphin	29-Jul-11	1	4	265	0	0
Rough-toothed dolphin	31-Jul-11	1	25	1507	3	1
Rough-toothed dolphin	31-Jul-11	2	12	540	0	0
Rough-toothed dolphin	31-Jul-11	3	7	262	0	0
Rough-toothed dolphin	01-Aug-11	2	9	186	1	0

Table 1 continued

Species	Date	Enc #	Group size	# photos	# genetic samples	# tag deployments
Rough-toothed dolphin	01-Aug-11	3	18	963	0	0
Rough-toothed dolphin	02-Aug-11	1	12	184	1	0
Rough-toothed dolphin	02-Aug-11	2	3	66	0	0
Rough-toothed dolphin	02-Aug-11	3	1	0	0	0
Rough-toothed dolphin	02-Aug-11	4	11	87	1	0
Rough-toothed dolphin	02-Aug-11	5	2	147	1	0
Rough-toothed dolphin	02-Aug-11	6	10	1156	3	2
Rough-toothed dolphin	02-Aug-11	7	4	47	0	0
Rough-toothed dolphin	06-Aug-11	2	7	503	0	0
Rough-toothed dolphin	08-Aug-11	2	14	727	2	0
Spinner dolphin	21-Jul-11	6	15	153	0	0
Spinner dolphin	22-Jul-11	2	75	1135	0	0
Spinner dolphin	25-Jul-11	5	30	0	0	0
Spinner dolphin	01-Aug-11	1	15	3	0	0
Spinner dolphin	01-Aug-11	4	4	42	0	0
Spinner dolphin	02-Aug-11	8	14	214	3	0
Spinner dolphin	03-Aug-11	2	2	0	0	0
Spinner dolphin	03-Aug-11	4	3	0	0	0
Spinner dolphin	04-Aug-11	1	80	1004	2	0
Spinner dolphin	04-Aug-11	3	35	158	0	0
Spinner dolphin	05-Aug-11	1	7	0	0	0
Spinner dolphin	05-Aug-11	3	70	791	0	0
Spinner dolphin	05-Aug-11	5	8	0	0	0
Spinner dolphin	06-Aug-11	3	4	0	0	0
Spinner dolphin	08-Aug-11	3	65	588	0	0
Unidentified dolphin	08-Aug-11	1	1	0	0	0
Unidentified odontocete	26-Jul-11	2	1	0	0	0
Unidentified odontocete	28-Jul-11	3	1	0	0	0
Unidentified odontocete	03-Aug-11	3	2	0	0	0

Table 2. Information on satellite tag deployments during July/August 2011¹.

Species	Tag ID	Date tagged	Individual ID	Sex ²	Duration of signal (days)	Tagged individual previously documented
Rough-toothed dolphin	SbTag001	31-Jul-11	HISb0860	Unknown	7.6	Yes
Rough-toothed dolphin	SbTag002	2-Aug-11	HISb0908	Male (g)	12.5	No
Rough-toothed dolphin	SbTag003	2-Aug-11	HISb0785	Male (g)	18.5	Yes
Bottlenose dolphin	TtTag002	3-Aug-11	HITt0632	Male (m)	34.1	No

¹All tagged individuals categorized as adults based on body size. ²Sex determined genetically (g) or by ventral morphology (m)

Table 3. Information on movements and habitats used by tagged individuals during July/August 2011.

Tag ID	# locations after filtering	Minimum cumulative distance moved (km)	Distance to deployment location (km) Median (Range)	Depth (m) Median (Range)	Distance from shore (km) Median (Range)
SbTag001	106	573.6	13.9 (2.3-39.7)	816 (318-2,162)	11.6 (3.7-19.5)
SbTag002	167	809.8	10.3 (0.6-30.2)	1,031 (495-2,762)	12.2 (5.4-21.5)
SbTag003	244	1,295.8	10.7 (0.9-42.4)	1,107 (383-2,703)	12.2 (3.6-21.7)
TtTag002	350	2,016.6	40.3 (0.2-61.4)	82 (2-2,144)	2.24 (0.12-12.93)




Figure 1. Map showing vessel tracklines and sighting locations from field efforts in July/August 2011.



Figure 2. Satellite-tagged rough-toothed dolphin, August 2, 2011. Photo © Robin W. Baird. This individual (HISb0785 in our catalog) has an unusual (piebald) coloration pattern, and had been previously documented off Kaua'i in October and November 2007 and in June 2008.




Figure 3. Satellite-derived locations of three tagged rough-toothed dolphins after filtering, during July and August 2011. See Table 2 for details.




Figure 4. Distance from deployment location over time for three rough-toothed dolphins satellite tagged in July and August 2011. All three individuals returned to the vicinity of the tagging location during the periods when locations were received.




Figure 5. Distance between pairs of tagged individual rough-toothed dolphins during the periods of tag overlap, during satellite overpasses when high-quality locations (LC3, LC2, LC1 only) were received from both tags. The similarity in appearance between the top and middle panel reflect that SbTag002 and SbTag003 remained in association for the period of tag overlap with SbTag001. Median distance apart: SbTag001 and SbTag002 = 13.5 km (max = 35.1 km); SbTag001 and SbTag003 = 13.7 km (max = 33.4 km); SbTag002 and SbTag003 = 1.06 km (max = 11.7 km). The x- and y-axis scales are the same for comparison.




Figure 6. Main cluster of a social network of distinctive and very distinctive rough-toothed dolphins photo-identified off Kaua'i and Ni'ihau from 2003 through 2011, with tagged individuals identified. All identifications of rough-toothed dolphins identified in 2003 and 2005 (see Baird et al. 2008) are included, with partial identifications of those photographed in 2008 and 2011. Each point represents an individual dolphin with lines connecting points for those individuals seen together in an encounter. Diagram produced with Netdraw 2.097 using a spring embedding layout. See Table 2 for tag numbers corresponding to individual identification numbers.




Figure 7. Locations of a satellite-tagged bottlenose dolphin from August 3, 2011 to September 6, 2011.