



# **Occurrence and Distribution of Rice's Whale Calls near De Soto Canyon, Gulf of Mexico 2022**

Melissa S. Soldevilla<sup>1</sup>, Amanda J. Debich<sup>2</sup>, Ludovic Tenorio<sup>2</sup>, Lance M. Garrison<sup>1</sup>

<sup>1</sup> NOAA Southeast Fisheries Science Center

<sup>2</sup> Cooperative Institute for Marine and Atmospheric Studies  
University of Miami Rosenstiel School of Marine and Atmospheric Science



NMFS Permit No. 779-1633-00

# Table of Contents

Executive Summary .....	3
Project Background.....	5
Rice’s Whales .....	7
Rice’s Whale Calls.....	7
Methods .....	9
Acoustic Recording Instrumentation .....	9
Data Collected.....	9
Data Analysis .....	10
Low Frequency Ambient Soundscape .....	10
Rice’s Whale Calls.....	10
Results and Accomplishments .....	12
Manuscript Preparation .....	<b>Error! Bookmark not defined.</b>
Moored Array Data Collection and Analyses .....	12
Low Frequency Ambient Soundscape .....	15
Rice’s Whale Long-moan Calls .....	22
Rice’s Whale Downsweep Sequence Calls.....	31
Vemco Acoustic Tag Detections.....	38
References.....	42

**This project is supported by funding from the Naval Sea Systems Command (NAVSEA) as part of the U.S. Navy’s marine species monitoring program.**

## Executive Summary

The Rice's whale (*Balaenoptera ricei*; formerly Gulf of Mexico (GOM) Bryde's whale) is estimated to have a population size of 51 individuals in U.S. waters (Garrison et al 2020) and was listed as endangered under the ESA in 2019 (84 FR 15446; 87 FR 8981). The majority of modern sightings occur in waters between the 100 – 400 m water depths in an area near the De Soto Canyon off northwestern Florida (Soldevilla et al. 2017, Rosel et al 2021). This primary distribution area is defined as the Rice's whale core habitat (Rosel and Garrison, 2022). Occurrence patterns from long-term passive acoustic monitoring (PAM) over the 2010-2018 period and from summer and fall visual surveys during 2018 and 2019 indicate the whales are found year-round within the core habitat, but also suggest there may be seasonal movements throughout, and potentially out of, this area. High densities of anthropogenic activities occur throughout the GOM, including oil and gas exploration and extraction, fisheries, shipping, and military activities. Many of these activities, including US Navy readiness training and testing, and Eglin Air Force Base activities, overlap with the whales' core habitat. Understanding seasonal distribution and density of Rice's whales throughout the core habitat will improve understanding of potential impact of human activities in this area, improve the accuracy and precision of impact assessments, and assist in developing effective mitigation measures as needed.

To improve management of human-based activities in the core habitat of these endangered whales, the Southeast Fisheries Science Center (SEFSC) began deploying a sparse array of 17 PAM units concurrent with one long-term HARP in May 2021. The PAM moorings are being deployed in two lines of 9 units each to nearly completely cover the core habitat over a nearly 2-year period to improve understanding of seasonal and interannual distribution, movement patterns, and habitat use. The moorings use SoundTrap ST500 and ST600 STDs, calibrated long-term recorders capable of continuously recording underwater sound in the 20 hertz (Hz) to 48 kilohertz (kHz) frequency range, including Rice's whale calls and ambient noise, for up to 6 months. Additionally, the study leverages a long-term HARP being deployed by the SEFSC, Scripps Institution of Oceanography, and collaborators, at the De Soto Canyon (DC) site in the core Rice's whale habitat over the August 2020 to July 2025 period. At this site, the team has been continuously recording ambient noise and other acoustic events in the 10 Hz to 100 kHz frequency range since 2010 to monitor the impacts of the Deepwater Horizon oil spill and subsequent restoration activities on cetaceans. Together with the sparse array of SoundTraps, these PAM deployments provide the necessary data to understand seasonal distribution and density of Rice's whales.

During 2022, work focused on continuing the 1.5 years of passive acoustic monitoring in the Rice's whale core habitat, including: 1) conducting fieldwork to service the array of PAM units to collect data recordings and continue data collection, and 2) conducting analyses of Rice's whale call presence in recordings from two 5-month deployments. In April 2022 and September 2022, SEFSC serviced the ST500 & ST600 STD SoundTraps at 17 mooring sites in the northeastern Rice's whale habitat. Recordings from the November 2021 to April 2022 deployment were recovered successfully from 16 of the sites, with instruments recording for a median of 4.4 months each, and most recordings ending in late March to early April 2022. Recordings from the April 2022 to September 2022 deployment were recovered successfully from 14 of the sites, with instruments recording a median of 4.9 months each, and most recordings ending in September 2022. Additional funds were leveraged from NOAA's Office of Protected Resources to complete one final deployment of the PAM array, recently recovered in March 2023, yielding close to two years of near-continuous recordings across sites over the May 2021 to March 2023 period. The concurrently deployed DC HARP will have three deployments spanning this period: 1) August 2020 to August 2021; 2) August 2021 to July 2022; and 3) July 2022 to July 2023. Data from the first and second HARP deployment have been recovered, yielding 13 months of concurrent data from 1 May 2021 to 3 June 2022.

During 2022–2023, data analyses were begun on the SoundTrap recordings (November 2021 to September 2022) as well as the concurrently deployed DC HARP recordings (August 2021 to June 2022). Automated spectrogram cross-correlation detectors for the downsweep-sequence and long-moan calls, developed under an earlier phase of this work, were run on all recordings. Given the critically endangered status of this species, automated detector thresholds are intentionally set low to minimize missed detections at the cost of increased false positive detections, and a subsequent manual validation step is conducted to remove false positive detections. This semi-automated process is both more efficient and consistent than a complete manual detection process and more accurate than a fully automated process. Across the 16 moorings recovered from the November 2021 to April 2022 period, there were a total of 1,835 instrument-days of recordings, a total of 533,193 Rice’s whale long-moan calls detected, and a total of 67,712 Rice’s whale downsweep sequences detected. The validation process has been completed for all long-moan and downsweep sequence calls from the 15 moorings with recordings available, yielding a total of 250,114 true long-moan call detections and 10,989 true downsweep sequence detections. During the November 2021 to April 2022 deployment period, true detections of Rice’s whale long-moans occurred at 14 sites, ranging from 368 to 53,884 calls per site. True detections of Rice’s whale downsweep sequences occurred at 10 sites, ranging from 18 to 3,148 calls per site. Similar to the May to September 2021 deployment data, higher numbers of detections occurred at sites on the inshore line than on the offshore line. Manual validation results indicate false detection rates for the long-moan detector and downsweep detector vary by site and over time within sites, with higher false-positive rates at offshore sites compared to inshore sites, and at the more southern sites near the Tampa shipping lane. High levels of seismic airgun activity during this deployment led to higher false-positive rates. Across sites, the daily occurrence of Rice’s whale long-moan calls varied by site as well, with long-moan calls present on 29 to 97 percent of days per site and downsweep sequence calls present on 0 to 62 percent of days per site over the Nov 2021 to April 2022 deployment period.


Across the 14 moorings recovered from the April to September 2022 deployment period, there were a total of 1,489 instrument-days of recordings that included 272,461 long-moan and 82,614 downsweep sequences detected by the automated detectors. Data storage hard-drives purchased during periods of supply chain issues caused by the COVID-19 pandemic have had a high rate of failure. Corrupted data were successfully recovered from 13 of the 14 sites, and the decompression, decimation, and automated detection processing steps have been completed.

Planned work for the remainder of the project includes completing validations and statistical analyses from the three deployments. The manual data validation process for the third SoundTrap deployment and concurrent HARP data are underway. Additionally, ambient noise analyses, monthly occurrence mapping, and evaluation of diel and seasonal changes in call occurrence and ambient noise impacts on call detection will be conducted. A manuscript will be drafted following completion of data analyses from the NOAA-funded fourth deployment. Finally, data collected during this project are being leveraged under NOAA-funded projects to acoustically track calling Rice’s whales throughout the core habitat and to evaluate feasibility of using spatially-explicit capture-recapture methods for density estimation.

## Project Background

The NOAA's Southeast Fisheries Science Center (SEFSC) and Scripps Institution of Oceanography (Scripps) have been collaboratively deploying long-term passive acoustic monitoring stations throughout the Gulf of Mexico (GOM) since 2010 to monitor the impacts of the Deepwater Horizon oil spill and subsequent restoration activities on cetaceans. High-frequency Acoustic Recording Packages (HARPs) deployed at the De Soto Canyon (DC) site in the Rice's whale core habitat have been continuously recording ambient noise and other acoustic events in the 10 Hz to 100 kHz frequency range. During 2019-2021, SEFSC conducted analyses of eight years of near-continuous DC HARP recordings (2010-2018) to understand Rice's whale seasonal and interannual occurrence patterns at this site. During the first phase of this project SEFSC developed automated Rice's whale call detectors and analyzed eight years of historic data from the DC HARP in the core habitat to establish complete occurrence time-series for understanding seasonal and interannual trends and for future habitat modeling and density estimation. Rice's whale call detections occurred throughout the year at this site with increased call detection rates during summer and fall compared to winter and spring.

In May 2021, the SEFSC implemented the second phase of this project to collect and analyze new passive acoustic data for at least one year over the entire Rice's whale core habitat to understand seasonal distribution patterns, density, and whale movements throughout the core habitat. To achieve this goal, the SEFSC developed a survey design using an array of 17 moored SoundTrap acoustic recorders deployed concurrent with the long-term DC HARP, with two lines of nine moorings each that nearly cover the Rice's whale core habitat (**Figure 1**). The SoundTrap mooring survey design included three 5-month deployment periods to collect over a full year of recordings at each site. Analytical objectives include running the automated long-moan and downsweep-sequence detectors developed in phase 1 on all recordings, with thresholds set to minimize missed detections at the cost of increased false positives, then conducting a manual verification step to remove all false positive detections and improve accuracy of the final results. Products to be developed include time-series of daily presence and total call detections by call type and site, time-series of ambient noise levels per site, and monthly maps of call detection rates and daily presence per site. These products of seasonal distribution and density of Rice's whales throughout the core habitat are needed to improve understanding of potential impact of human activities in this area, improve the accuracy and precision of impact assessments, and assist in developing effective mitigation measures as needed.



**Figure 1.** Historic long-term passive acoustic monitoring station (HARP; dark blue) deployed in the Rice's whale core habitat in the northeastern Gulf of Mexico since 2010, and sparse array passive acoustic monitoring stations (SoundTraps; light blue) deployed over the 2021-2023 period. Circles around passive acoustic stations indicate the expected acoustic coverage, assuming 20 km call detection distances. The NMFS core habitat for Rice's whales is indicated as a shaded polygon. The long-term De Soto Canyon (DC) HARP site, where Rice's whale calls have previously been detected, is being deployed concurrent with the SoundTrap array under a Deepwater Horizon Restoration project.

## Rice's Whales

The Rice's whale (*Balaenoptera ricei*; formerly Gulf of Mexico Bryde's whale), estimated to have a population size of 51 individuals in US waters (CV 0.53, Garrison *et al.*, 2020), was listed as endangered under the US Endangered Species Act (ESA) in 2019. The majority of modern sightings occur in waters between the 100 – 400 meter (m) water depths near the De Soto Canyon off northwestern Florida (Soldevilla *et al.*, 2017; Rosel *et al.*, 2021), an area defined as the Rice's whale core habitat (Rosel & Garrison, 2022). Occurrence patterns from long-term passive acoustic monitoring over the 2010-2018 period and two recent summer and fall surveys in 2018-2019 indicate the whales are found year-round within this core habitat. Results also show decreased call detection rates in winter and spring, indicating there may be seasonal movements throughout the core habitat, and potentially beyond it to areas like the recently identified habitat along the shelf break off Louisiana (Soldevilla *et al.*, 2022a). High densities of anthropogenic activities occur throughout the GOM, including oil and gas exploration and extraction, fisheries, shipping, and military activities and several of these activities overlap with the whales' core habitat. Many of these activities, including US Navy readiness training and testing and Eglin Air Force Base activities, overlap with the whales' core habitat. Understanding seasonal distribution and density will improve understanding of potential impact of human activities in the core habitat, improve the accuracy and precision of impact assessments, and assist in developing effective mitigation measures as needed.

## Rice's Whale Calls


Long-term, broad-coverage passive acoustic monitoring is a highly effective tool for investigating whale seasonal and interannual occurrence patterns. In the GOM, three call types have been identified and definitively attributed to free-ranging Rice's whales (Rice *et al.*, 2014, Širović *et al.*, 2014, Soldevilla *et al.*, 2022b) and one additional call type has been proposed as a likely candidate (Širović *et al.*, 2014; **Figure 2**).

### ***Downsweep Pulse Calls***

Rice's whales produce downsweep pulse sequence calls made up of series of two or more short-duration downsweeps (mean: 8 downsweeps, range: 2-25) ranging from  $110 \pm 4$  to  $78 \pm 7$  Hz, with a mean duration of  $0.4 \pm 0.1$  s, an inter-pulse interval of  $1.3 \pm 0.1$  s, and source levels of  $155 \pm 14$  dB re:  $1 \mu\text{Pa}$  at 1 m (Rice *et al.*, 2014, Širović *et al.*, 2014, Soldevilla *et al.*, 2022b). A second downsweep call type, higher in frequency (170 to 110 Hz), segmented, and typically occurring in repeated sequences of doublets, also has been detected in autonomous recordings and is proposed to be a possible Rice's whale call (Širović *et al.*, 2014).

### ***Tonal Calls***

Rice's whales produce two tonal call types: long-moan calls and tonal-sequence calls (Rice *et al.*, 2014, Soldevilla *et al.*, 2022b). The long-moan call type is a long-duration, amplitude-modulated downsweep ranging from 150 to 75 Hz with a mean center frequency of 107 Hz, mean 22.2 s duration, and 3.4 pulse/s amplitude pulse rate (Rice *et al.*, 2014, Soldevilla *et al.*, 2022b). Stereotyped variants of the long moan that are common in the western Gulf are occasionally detected in the core habitat as well (Soldevilla *et al.*, 2022a). The second tonal call type, the tonal-sequence, consists of 1-6 narrow-band constant-frequency tones in sequence following some long-moans, with individual tonals having a mean center frequency of 103 Hz and mean 3.6 s duration (Rice *et al.*, 2014).



**Figure 2.** Spectrograms of Rice's whale calls and potential calls

## Methods

### Acoustic Recording Instrumentation

#### *High-frequency Acoustic Recording Package (HARP)*

HARPs were used to record marine mammal sounds and characterize the low-frequency ambient soundscape in the GOM at the DC HARP site from 2010 through 2023. HARPs can autonomously record underwater sounds from 10 Hz up to 160 kHz and are capable of approximately 300 days of continuous data storage. The HARPs were deployed in either a seafloor mooring or a seafloor package configuration with the hydrophones suspended 10 m above the seafloor (**Figure 4**). Each HARP is calibrated in the laboratory to provide a quantitative analysis of the received sound field. Representative data loggers and hydrophones were also calibrated at the Navy's TRANSDEC facility to verify the laboratory calibrations (Wiggins & Hildebrand 2007).


#### *SoundTrap ST500 & ST600 (SoundTrap)*

SoundTrap ST500 and ST600 STD recorders (Ocean Instruments Inc.) were deployed as a sparse array to record marine mammal sounds and characterize the low-frequency ambient soundscape throughout the Rice's whale core habitat in the northeastern GOM over the May 2021 to September 2022 period. The SoundTrap ST500 and ST600 STDs are calibrated long-term recorders capable of continuously recording underwater sound in the 20 Hz – 60 kHz frequency range, including Rice's whale calls and ambient noise, for up to six months. The SoundTraps were deployed in a small mooring configuration with the hydrophones suspended 3 m above the seafloor. The ST500 & ST600 STD recorders are factory calibrated at 250 Hz. The SoundTrap moorings use a Vemco VR2AR acoustic release that allows opportunistic collection of transmissions from Vemco-acoustic-tagged fish and reptiles that pass by the mooring.

### Data Collected

Data were collected by SEFSC and Scripps from the historic DC HARP site (29° 2.878' N 86° 05.847' W, 270 m depth) during the August 2020 - 2023 period using HARPs sampling at 200 kHz, under funding from a *Deepwater Horizon* Restoration project. The DC HARP site (DCH) is located approximately in the center of the Rice's whale core habitat area (**Figure 1**; Rosel & Garrison, 2022).

Concurrent data were collected by SEFSC from a sparse array of 14 SoundTrap moorings deployed in two lines that nearly completely covered the core habitat (**Figure 1**) during three deployments over the May 2021 to September 2022 period. A fourth deployment, funded by NOAA Office of Protected Resources (OPR), collected data over the September 2022 to March 2023 period. The SoundTraps sampled at 24 kHz over the periods: 1) May to October 2021; 2) November 2021 to April 2022; 3) May to September 2022; and 4) September 2022 to March 2023. The concurrently deployed DC HARP sampled over the August 2020 to August 2021 and August 2021 to July 2022 periods and is currently collecting data over the July 2022 to July 2023 period.



**Figure 3.** Schematic of a HARP seafloor package

## **Data Analysis**

Recording over a broad frequency range of 10 Hz to 100 kHz allows detection of the low-frequency ambient soundscape, baleen whales (mysticetes), toothed whales (odontocetes), and anthropogenic sounds. Because analyses were focused on the Rice's whale and ambient noise, only the low-frequency data were required for these analyses. The HARP recordings were decimated by a factor of 100 and the SoundTrap recordings were decimated by a factor of 12 to provide an effective bandwidth of 10 Hz to 1 kHz. Long-term spectral averages (LTSAs) were created from the decimated data with a 1 Hz frequency and 5 s temporal resolution.

### **Low Frequency Ambient Soundscape**

All recordings were converted to sound pressure levels using factory calibration values for SoundTrap recordings and calibration values obtained from full-system calibrations conducted at the U.S. Navy's Transducer Evaluation Center in San Diego, CA for HARP recordings. Hourly spectral averages and associated standard deviations were computed by combining sound pressure spectrum levels calculated from each acoustic record per hour. System self-noise was excluded from these averages. Time series of the 1, 50, and 99 percentiles of the average hourly spectrum levels at 100 Hz and 125 Hz were developed from these data. They were also combined to obtain monthly spectral averages to evaluate longer-term changes in the ambient soundscape and its potential impacts on baleen whale call detectability.


### **Rice's Whale Calls**

#### ***Automated Call Detectors***

During prior work conducted in 2018-2019, spectrogram cross-correlation detectors for long-moan calls and downsweep pulse sequences were developed in Ishmael (Mellinger & Clark 2000) using a two-day training dataset and a separate testing dataset to characterize miss rates and false detection rates. In 2021, these detectors were run on spectrograms of recordings from all SoundTrap sites and the concurrent DC HARP recordings from the 1<sup>st</sup> deployment covering the May 2021 to November 2021 period. In 2022, the detectors were run on spectrograms of recordings from all SoundTraps and HARPs deployed over the 2<sup>nd</sup> and 3<sup>rd</sup> deployments covering the November 2021 to September 2022 period. For all analyses, spectrograms were calculated using an FFT frame size of 512 samples, no zero-padding, 50% overlap, and spectrogram equalization with 3 s spectral averaging.

#### ***Long-Moan Detector Settings***

Long-moan call contours contain five sections which include the preliminary upsweep, the approximately 150 Hz tone, the first part of the downsweep (slope 1), the second part of the downsweep (slope 2), and the long nearly-constant-frequency tail (**Figure 5**). The cross-correlation contour kernel for the long-moan call focused on the 150 Hz tone and slope 1, the most consistent parts of the frequency-modulated tonal call. The kernel contour is defined by a 1.1 s tone from 146 Hz to 145 Hz followed by a 3.7 s downsweep from 145 Hz to 112 Hz, each with a 14 Hz contour bandwidth. Detection function smoothing was enabled. The detection threshold was set to 4.5, and minimum and maximum detection durations were 0.5 s and 3.0 s, respectively. The minimum time allowed between subsequent detection events was 0.5 s. The threshold of 4.5, yielding a 6.4% missed call rate and a 26.4% false detection rate on a test dataset, was selected to minimize miss rates without excessive false detection rates. Missed detections were typically associated with calls with low signal to noise ratios. The majority of false alarms were associated with disk write noise from the recording instrument and tonal sounds from passing ships.



**Figure 4.** Five sections of a long moan call. Two sections, the 150 Hz tone and slope 1 were used to create the contours in the long-moan detector.

### ***Downsweep Pulse Sequence Detector Settings***

The *Ishmael* downsweep pulse sequence detector used the regular sequence feature to detect sequences of individual downsweep pulses as a single call. The cross-correlation contour kernel was defined as a single 4 s downsweep from 120 Hz to 80 Hz, with a 20 Hz contour bandwidth. For regular sequences, the minimum and maximum repetition period between individual pulse detections were set to 0.9 s and 1.1 s, respectively, and an 11 s window with 75% overlap was used. The detection threshold was set to 11, and minimum and maximum detection durations were set to 0.1 s and 40 s, respectively. The minimum time allowed between detection events was 0.4 s. The threshold of 11, yielding a 12.6% missed sequences rate and a 69.1% false detection rate on a test dataset, was selected to minimize miss rates without excessive false detection rates. Missed detections were typically associated with calls with low signal to noise ratios. The majority of false alarms were associated with long-moan calls with strongly pulsed tails and seismic survey airgun pulses with unusually short inter-pulse intervals or strong multipath effects.

### ***Validation of automated call detections***

Given the critically endangered status of Rice's whales, automated detector thresholds were intentionally set to minimize missed detections at the cost of increased false positive detections, rather than selecting a threshold with equal miss and false alarm rates. The threshold selections aimed to reduce missed detections as much as possible while balancing the need to keep false detections within a reasonable number. Therefore, these preliminary detections require a follow-up step to manually validate and remove all false detections for a final dataset. This semi-automated process is both more efficient and consistent than a complete manual detection process and more accurate than a fully automated process. In the validation step, each automated detection is manually reviewed and scored as a true or false detection, and false positive rates are calculated as the percentage of false positives to total detections. In 2021, all detections were manually validated for long-moan call detections at 12 of the 15 DC array sites from the first deployment over the May to September 2021 period. In 2022, manual validation of long-moan detections at the remaining 3 sites, and manual validation of downsweep sequence detections at all 15 sites were completed. Additionally, in 2022, manual validation of long-moan and downsweep sequence detections was completed for all 15 SoundTrap sites from the second deployment over the November 2021 to April 2022 period. Only the automated detector has been run for Soundtrap recordings from the third deployment from the April to September 2022 period and for HARP recordings from the August 2021 to June 2022 period; manual validation of long-moan and downsweep sequence detections is planned.

## Results and Accomplishments

### Moored Array Data Collection and Analyses

In April 2022, the SEFSC recovered 16 of 17 SoundTraps from the second deployment (November 2021 to April 2022) and deployed 16 SoundTraps for the third deployment from NOAA's *R/V Southern Journey*. One unit (site DCO) was not recovered as the acoustic release would not communicate. In September 2022, the SEFSC recovered 14 of 16 SoundTraps from the third deployment and deployed a final round of 13 SoundTraps (funded by NOAA's Office of Protected Resources). Two units (sites DCF and DCR) from the third deployment were not recovered as the acoustic release communications were sparse and inconsistent despite substantial effort on site to improve communications for recovery. During this trip, the SEFSC additionally recovered and redeployed the HARP at site DCH. SoundTrap moorings from the fourth and final deployment were recovered in March 2023, and the HARP will be recovered in July 2023 as part of the *Deepwater Horizon* Restoration passive acoustic monitoring project.

The acoustic recordings successfully recovered from 16 of the 17 SoundTrap locations from deployment 2 yielded a total of 2,120 instrument-days (50,876 hours) of recordings over the November 2021 to April 2022 period (**Table 1**). The SoundTraps recorded for a median of 4.4 months each (range 0.3 to 4.7 months), with recordings ending between mid-February to early April (some were still recording on recovery) and yielded high quality recordings throughout the deployment at 14 of the 16 sites. Site DCB had power consumption issues and only recorded poor quality data for 5 days while the ST600 deployed at site DCE flooded and no quality recordings could be recovered. The SoundTraps deployed at sites DCG, DCI, and DCM had hydrophone malfunctions that led to impulsive noise occurring periodically through the recordings starting in February. Data from these three sites are undergoing QA/QC to remove low-quality data and retain useable recordings where possible.

The acoustic recordings successfully recovered from 14 of the 16 SoundTrap moorings from deployment 3 yielded a total of 1,489 instrument-days (35,724 hours) of recordings over the April to September 2022 period (**Table 1**). The SoundTraps recorded for a median of 4.9 months each (range 0.3 to 5.6 months), with recordings ending between June to mid-September (one ST600 was still recording on recovery) and yielded high quality recordings throughout the deployment at 11 of the 14 sites. Site DCL had power consumption issues and only recorded poor quality data for seven days (these could not be repaired following Serial Advanced Technology Attachment (SATA) hard drive corruption); Site DCQ had connection issues with the hydrophone, with signal dropouts during the first two weeks and no sound in recordings after two weeks; and the ST600 deployed at site DCN flooded and no recordings could be recovered. The SoundTraps deployed at sites DCC, DCD, DCG, and DCP appear to have had hydrophone malfunctions that led to impulsive noise occurring periodically through the recordings. Data from these four sites are undergoing QA/QC to remove low-quality data and retain useable recordings where possible. The HARP deployed concurrent with the 2<sup>nd</sup> and 3<sup>rd</sup> deployments of SoundTraps yielded 285 days (6,827 hours) of high-quality recordings over the August 2021 to June 2022 period.

As noted above, technical challenges continue with the SoundTraps, including the ST500s built in the early days of the COVID pandemic and the ST600s built in summer 2021. While some of ST500 problems are associated with the internal lithium batteries remaining in a fully discharged state for an extended period due to inaccessibility during mandatory COVID closures in 2020-2021, other challenges are associated with hydrophone failures; this model is no longer being manufactured by Ocean Instruments due to reliability issues. The newer model ST600s are generally more reliable with respect to hydrophone quality and power consumption; however, instrument flooding, experienced by many NOAA and academic users, has affected this project as well with two ST600 flooding events. In all instrument failure cases, the

SoundTraps exhibiting problematic behavior have been returned to the manufacturer and repaired prior to redeployment, except in a couple of cases when the failure was not evident until returning from sea. In addition to these challenges, SATA hard-drives were difficult to purchase during the pandemic when supply-chain issues for products manufactured in China were an issue. We have experienced a high number of failures with SATA drives purchased during this time; both the original and backup SATA data storage drives for deployment 3 were corrupted during the backup process. All corrupted drives were sent to repair companies and nearly all of the data were successfully recovered. A third drive containing the second half of the recordings also failed following decompression, decimation, and auto-detection processing but the original files were recovered from the network backup. These hard-drive failures caused delays to completing analyses of deployment 3, but everything is now back on track and moving forward as expected.

The preliminary results of acoustic data analysis from recordings at the 18 DC array sites deployed during the first 3 deployments over the May 2021 to September 2022 period are summarized below. At this time, LTSAs and daily sound pressure spectrum levels have been calculated for all recordings at the 18 sites, the two automated spectrogram cross-correlation detectors have been run on all recordings from the 18 sites, and the resulting long-moan and downsweep sequence detections have been manually validated for recordings from deployments 1 and 2 at all 17 SoundTrap sites. The validation of long-moan and downsweep detections from the third deployment will be completed in 2023. Here we describe the low-frequency ambient soundscape, hourly and daily Rice's whale call presence, daily call detections per site, and map the monthly distribution of call rates and daily occurrence per site. Final statistical analyses of spatial distribution and variation in Rice's whale call occurrence and ambient noise levels will be completed following the completion of data collection and analysis from the third and fourth deployments, during 2023.

**Table 1.** Acoustic monitoring effort at 18 sites near De Soto Canyon during three deployments over the May 2021 to September 2022 period.

<sup>1</sup> The HARP at site DCH began recording in August 2020; only data collected concurrent with the SoundTrap array are included in these analyses.

<sup>2</sup> These two SoundTrap moorings had hydrophone malfunctions during deployment 1 and parts of the recordings are corrupted.

<sup>3</sup> Successive mooring deployment locations occurred within 25m, except at sites A & P. Deployment 3 at Site A was moved 6.5km southeast to deeper waters at 29.5675 N, -87.3229 W, as the original site was too shallow (85m). Deployment 2 at Site P was moved 3.6km southeast to the originally planned site location at 28.0137 N, -84.9703 W, as it was misplaced during the first deployment.


Site	Latitude (°N)	Longitude (°W)	Deployment 1				Deployment 2				Deployment 3			
			Start Time	End Time	Effort (Days)	Effort (Hours)	Start Time	End Time	Effort (Days)	Effort (Hours)	Start Time	End Time	Effort (Days)	Effort (Hours)
DCA <sup>3</sup>	29.5640	-87.3815	5/2/21 00:58	9/28/21 23:57	150	3,599	11/17/21 13:02	4/2/22 00:13	135.5	3251	4/2/22 00:43	9/6/22 09:44	157.4	3777
DCB <sup>2</sup>	29.7579	-87.0380	5/2/21 03:32	9/10/21 18:30	131.6	3,159	11/11/21 13:00	-	-	-	4/10/22 17:53	9/9/22 17:35	152.0	3648
DCC	-	-	-	-	-	-	11/16/21 23:04	4/1/22 09:34	135.4	3250	4/10/22 15:57	7/5/22 22:07	86.3	2070
DCD	29.5626	-86.6830	5/2/21 06:05	8/24/21 03:43	113.9	2,734	11/11/21 16:58	3/27/22 05:47	135.5	3253	4/10/22 13:24	8/23/22 03:40	134.6	3230
DCE	29.2153	-86.7550	5/2/21 08:31	10/2/21 19:53	153.5	3,683	11/16/21 20:05	11/26/21 00:18	9.2	220	4/4/22 20:05	9/2/22 03:26	150.3	3607
DCF	29.3206	-86.3683	5/2/21 16:03	9/16/21 04:36	136.5	3,277	11/11/21 20:59	4/4/22 23:06	144.1	3458	4/4/22 23:00	-	-	-
DCG	28.9647	-86.4710	5/2/21 18:41	8/23/21 02:47	112.3	2,696	11/16/21 17:16	3/30/22 22:52	134.2	3222	4/4/22 16:53	7/18/22 13:51	104.9	2517
DCH <sup>1</sup>	29.0554	-86.0965	5/1/21 00:00	8/23/21 01:05	114	2,737	8/23/21 06:00	6/3/22 18:23	284.5	6828	-	-	-	-
DCI <sup>2</sup>	28.7269	-86.1753	5/2/21 21:19	5/19/21 10:50	16.6	398	11/16/21 14:24	3/10/22 00:16	113.4	2722	4/2/22 12:27	9/11/22 08:13	161.8	3884
DCJ	28.8284	-85.7759	5/2/21 23:41	9/14/21 07:41	134.3	3,224	11/14/21 16:33	3/31/22 10:32	136.7	3282	4/4/22 19:04	9/21/22 07:33	169.5	4068
DCK	28.4830	-85.8975	5/3/21 02:12	9/16/21 15:51	136.6	3,278	11/14/21 19:42	4/1/22 17:18	137.9	3310	4/2/22 14:58	8/31/22 15:14	151.0	3624
DCL	28.5542	-85.5043	5/3/21 04:52	9/22/21 19:53	142.6	3,423	11/14/21 23:30	4/3/22 23:50	140	3360	4/4/22 00:39	-	-	-
DCM	28.2201	-85.6354	5/3/21 11:14	9/23/21 02:07	142.6	3,423	11/15/21 15:30	2/18/22 20:00	95.2	2285	4/2/22 18:25	9/2/22 03:40	152.4	3657
DCN	28.2865	-85.2375	5/3/21 13:50	9/21/21 00:10	140.4	3,370	11/15/21 12:28	4/3/22 20:58	139.4	3345	4/3/22 21:46	-	-	-
DCO	27.9783	-85.3304	5/3/21 16:11	9/23/21 23:38	143.3	3,439	11/15/21 18:32	-	-	-	-	-	-	-
DCP <sup>3</sup>	28.0225	-85.0012	5/3/21 18:17	8/10/21 09:23	98.6	2,367	11/16/21 00:00	4/3/22 16:40	138.7	3329	4/3/22 18:39	6/1/22 11:27	58.7	1409
DCQ	-	-	-	-	-	-	11/16/21 04:20	2/28/22 09:54	104.2	2502	4/3/22 13:46	4/13/22 05:34	9.7	232
DCR	-	-	-	-	-	-	11/16/21 01:15	3/31/22 19:54	135.8	3259	4/3/22 14:53	-	-	-

## Low Frequency Ambient Soundscape


Across the three deployments, the low-frequency soundscape in the 10-800 Hz range showed strong similarities across most of the sites in the DC array. Long-term spectrograms of the full deployment at each site show that the 100-800 Hz band is primarily dominated by wind and wave noise with broadband noise level increases seen across all sites at the same time (**Figure 5**). Seismic airgun surveys, with strong energy in the 10-70 Hz band showing distinctive energy peaks, dominate the low frequency band during the 2<sup>nd</sup> and 3<sup>rd</sup> deployments, while they were only evident in the first three weeks of May 2021 and in late August and September 2021 of the 1<sup>st</sup> deployment (**Figure 5**). Shipping noise is ubiquitous across all sites in the 30-100 Hz band when airgun noise doesn't mask it, and is particularly strong at the eight southernmost sites (**Figure 5**). The recorder at site DCA was deployed in 85 m depths, on the shelf, for deployments 1 & 2, while all other sites were deeper (180-450m) and the soundscape at this site was distinctly different from other sites, with less seismic airgun noise, more shipping noise, and the presence of biological noise from fish, including diel chorusing (e.g. bands around 200 and 400 Hz; **Figure 5**). Long-term spectral averages also give a quick look into data quality, and the issues described above at sites DCB, DCG, DCI, and DCM during the 2<sup>nd</sup> deployment and sites DCC, DCD, DCG, DCP, and DCQ during the 3<sup>rd</sup> deployment are evident in these figures (**Figure 5**).

- While the 2021 Atlantic hurricane season was an unusually active one with seven named storms passing through the GOM during deployment 1, only one tropical storm (Tropical Storm Alex) passed through the Gulf during the 3<sup>rd</sup> deployment period in 2022. Similar to 2021, increases in broadband noise levels over the 100-1000 Hz band were evident across all sites during the storm on June 5-6 (**Figure 5**).
- Short-term increases in broadband noise levels in the 100-1000 Hz band associated with heavy weather were evident across sites during winter and spring, from Dec 2021 to May 2022 (**Figure 5**). During these periods, lower frequency noise levels associated with seismic surveys often decreased.
- Site DCA, the shallowest site at 85 m and the only site on the shelf, had a different soundscape than all the other sites (**Figure 5**). The soundscape was characterized by more shipping noise, less seismic survey noise, and high levels of biological activity including diel fish choruses. The soundscape only changed slightly at this site when moved to deeper waters (198 m) for deployment 3, when compared with noise levels during the same season from deployment 1.
- Similar to deployment 1 data from 2021, a comparison of sound pressure spectrum levels at site DCH deployment 2 (the fully calibrated HARP) and the SoundTrap sites from deployments 2 & 3 (factory calibrated at 250 Hz) shows that sound levels below 20 Hz drop off more rapidly on the SoundTraps than on the HARP, indicating that ambient noise quantification from SoundTraps is inappropriate at these lower frequencies (**Figure 5**).
- Seismic survey noise in the 10-70 Hz range occurred nearly constantly and at higher levels during the 2<sup>nd</sup> and 3<sup>rd</sup> deployments compared the infrequent occurrence during the 1<sup>st</sup> deployment (**Figure 5**). The same airgun surveys were detected at nearly all sites across the array from late August 2021 through September 2022, with brief gaps in activity evident in March 2022 at southern sites and June of 2022 across sites. Airgun survey noise was limited at northernmost sites A & B.
- Diel date vs time plots of noise levels at 125 Hz (the center frequency for Rice's whales long-moans that typically range from 150 to 100 Hz) show how 125 Hz noise levels vary by time of day and day of the year (**Figure 6**). The slight diel differences seen at sites DCB and DCD during the 1st deployment are not evident during the 2<sup>nd</sup> and 3<sup>rd</sup> deployments. A tidal signal is evident during the 3<sup>rd</sup> deployment at site DCE (**Figure 6**). This may be related to fish chorusing since tidal strumming is not apparent in the long-term spectrogram (**Figure 5**).


- Large scale weather events described from long-term spectrograms are also evident in the 125 Hz diel plots (**Figure 6**), e.g., during April through June 2022, but these are not as apparent as the tropical storms and hurricanes seen during the 1<sup>st</sup> deployment (**Figure 6**)
- Noise levels at 125 Hz were generally higher at the shelf site DCA and at the eight southern sites, particularly the southern inshore sites (**Figure 6**). This was especially evident during the 2<sup>nd</sup> deployment.
- Shorter duration events, seen as brighter yellow or red spots in diel plots, represent ship passings, and are seen across all deployments at all sites, in particular the 5 southern inshore sites and the shallow DCA site (**Figure 6**).
- Time-series of the daily statistical distribution of average hourly sound pressure spectrum levels at 125 Hz (**Figure 7**) follow similar patterns of increased levels at the time periods described above. Daily spectrum levels are generally higher and more variable at the more southern sites compared to northern sites, with the exception of site DCA. At many sites, median noise levels track closely with the lowest 1<sup>st</sup> percentile noise levels, particularly at the northern sites.
- Noise levels at 100 Hz (the center frequency for Rice's whales downsweep sequences that typically range from 75 to 125 Hz) largely follow the same temporal patterns as those seen at 125 Hz, but at higher levels and with ship passing events occurring for longer durations (**Figures 8 & 9**).




**Figure 5.** Hourly median long-term spectral averages of the SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022 showing recorded ambient noise levels from 10-1000 Hz. Gray indicates periods with no recording effort. Some dark yellow and red periods (e.g. sites B, C, G, I, and P) and dark blue periods (e.g. sites B, L, and Q) represent periods with poor quality data that are currently undergoing QA/QC to salvage any good quality recordings; poor quality recordings will be removed.




**Figure 6.** Diel variation in hourly median sound pressure levels at 125 Hz (center frequency of long-moan calls) for SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022. Gray indicates periods with no recording effort. Some dark yellow and red periods (e.g. sites B, C, G, I, and P) and dark blue periods (e.g. sites B, L, and Q) represent periods with poor quality data that are currently undergoing QA/QC to salvage any good quality recordings; poor quality recordings will be removed.



**Figure 7.** Daily distribution (light gray shading: 1 – 99 percentile; gray line: 50 percentile) of average hourly sound pressure levels at 125 Hz (center frequency of long-moan calls) for SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022. Dark gray indicates periods with no recording effort. Some periods with potential poor quality recordings are currently undergoing QA/QC to salvage any good quality recordings; poor quality recordings will be removed.



**Figure 8.** Diel variation in hourly median sound pressure levels at 100 Hz (center frequency of downsweep sequence calls) for SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022. Gray indicates periods with no recording effort. Some dark yellow and red periods (e.g. sites B, C, G, I, and P) and dark blue periods (e.g. sites B, L, and Q) represent periods with poor quality data that are currently undergoing QA/QC to salvage any good quality recordings; poor quality recordings will be removed.



**Figure 9.** Daily distribution (light gray shading: 1 – 99 percentile; gray line: 50 percentile) of average hourly sound pressure levels at 100 Hz (center frequency of downsweep sequence calls) for SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022. Dark gray indicates periods with no recording effort. Some periods with potential poor quality recordings are currently undergoing QA/QC to salvage any good quality recordings; poor quality recordings will be removed.

## Rice's Whale Long-moan Calls

There were a total of 1,316,461 automated detections of Rice's whale long-moan calls in recordings from the 18 sites in the DC array during the May 1, 2021 – September 21, 2022 period, ranging between 434 and 85,354 detections per deployment per SoundTrap site (**Table 2**). A total of 144,819 detections occurred on the 2<sup>nd</sup> HARP deployment at DCH, which spans both the 2<sup>nd</sup> and 3<sup>rd</sup> deployments of SoundTraps. Over the 1<sup>st</sup> and 2<sup>nd</sup> deployments at sites for which automated long-moan detections have been manually validated, 496,667 of the 899,181 detections were validated as true long-moan calls, with a range of 7 to 53,884 per deployment (**Table 2**). True long-moan call detections were higher at inshore sites than offshore across both deployments (**Table 2, Figures 10, 11**). False detection rates per deployment averaged 32.6% and 53.1% for the 1<sup>st</sup> and 2<sup>nd</sup> deployments respectively, and ranged between 13.3% and 100%. The sites with higher false positive rates (>50%) were the offshore sites where Rice's whale calls were less common (some of these will be reduced following removal of poor quality data), and the southernmost inshore site DCR where vessel noise from the Tampa shipping fairway was common (**Table 2, Figure 11**). False detection rates also varied over time within sites, with increased rates at sites in the southern half of the array from September 2021 to March 2022, coincident with increased activity from seismic airgun surveys, which were the cause of most of the false detections (**Figure 11**).


Preliminary results indicate true long-moan calls were present during an average of 78% of days (range 7-99% of days) per site and during an average of 46% of hours (range 0.5-87% of hours) with recording effort per site during the two deployments from May 2021 to April 2022 deployment (**Table 2**). Daily and hourly presence of Rice's whale calls were higher at inshore sites (mean: 89% of days, range 61-99%; mean 67% of hours, range 37-87%) than offshore sites (mean: 59% of days, range 7-93%; mean 26% of hours, range 0.5-62%). A preliminary comparison with potential detection ranges (**Figure 12**) as a function of noise levels at 125 Hz (e.g. **Figure 7**) indicates that masking effects of noise levels are not the primary driver for this inshore/offshore difference as noise levels are generally lower offshore leading to greater estimated detection ranges at offshore sites compared to inshore sites (future analyses will include site-specific sound propagation conditions). Preliminary results suggest potential seasonal movements between southern and northern sites as call rates and daily call presence vary across sites over time (**Figures 11, 13, 14, 15**) with higher occurrence and call detection rates at northern sites (DCB, DCD, DCF) in May through July 2021 compared to later months, and higher occurrence and call detection rates at southern sites (DCJ, DCL, DCN, DCP) during August 2021 through February 2022. Spatio-temporal variation will be re-evaluated upon completing the validation at the DCH site and data from the 3<sup>rd</sup> and 4<sup>th</sup> array deployments.

- The spectrogram cross-correlation detector for long-moan calls yielded a total of 1,316,461 detections in recordings from the 18 sites over the three deployments from May 2021 to September 2022, and ranged between 434 and 85,354 per SoundTrap site (**Table 2**).
- Validation of auto-detections of long-moan calls has been completed on the remaining three of 15 sites from the 1<sup>st</sup> deployment and on all 15 SoundTrap sites from the 2<sup>nd</sup> deployment. True long-moan calls were detected at every site with quality recordings, with higher numbers of detections per deployment at inshore sites (mean: 28,838, range: 10,208 to 53,884) compared to offshore sites (mean: 4,273, range: 7 – 11,650; **Table 2, Figures 10, 11, 13**).
- Across all validated detections from the 1<sup>st</sup> and 2<sup>nd</sup> deployment, no western long-moan variants were detected at any site during the May 2021 – April 2022 period.
- Validation results from the 18 sites yielded false detection rates averaging 32.6% and 56.4% for the 1<sup>st</sup> and 2<sup>nd</sup> deployments, respectively, and ranging between 13.3% and 100% for the long-moan call detector per deployment across the 18 sites over the May 2021– April 2022 period.


- The highest false detection rates (>90%) occurred during both deployments at site DCA, and during deployment 1 at DCI and deployment 2 at DCE and DCQ. At DCA, the mooring was deployed in 85 m water depth where Rice's whale calls were rare while fish chorusing and vessel noise were common. Sites DCE and DCI had hydrophone malfunctions, with DCE having no true detection while DCI was unreliable for the 17 days of recordings, though some true long-moan calls were still detected. High false positive rates (93%) at site DCQ occurred when seismic survey noise was prevalent, with concurrently high false positive rates at site DCR (73%) just inshore of it and DCP (58%), one site to the north and inshore (**Figure 11**).
- Beyond these, false detection rates were higher at offshore sites (30-84%), where fewer true calls were detected, compared to inshore sites (13-40%). False detections also varied over time within sites, with increased rates at sites in the southern half of the array from September 2021 to March 2022, coincident with increased activity from seismic airgun surveys, which were the cause of most of the false detections (**Figure 11**). A similar effect was seen at site DCO in May.
- Preliminary results indicate that true Rice's whale long-moan calls were present an average of 78% (range: 7 – 99%) of days per site, and were present in an average of 46% (range: 0.5 to 87%) of recording hours per site during the two deployments from May 2021 to April 2022 deployment (**Table 2**). Percent of days and percent of hours present were generally higher at the inshore sites (mean 89%, range 61-99% of days; mean 67%, range 37-87% of hours) than the offshore sites (mean 60%, range 7 – 93% of days; mean 26%, range 0.5 – 62% of hours).
- Preliminary results of estimated maximum detection ranges of long-moan calls as a function of ambient noise conditions at 125 Hz (assuming call source level of 145 dB, detection threshold of 10 dB SNR, and 17logR geometric spreading loss at all sites) suggest detection ranges generally are higher at offshore sites where noise levels are lower. This suggests noise masking is not the cause of lower detection rates at offshore sites. Future analyses will include sound propagation modeling using parabolic equations to better estimate site-specific transmission loss effects on detection ranges (**Figure 12**).
- At the southern sites (DCK to DCR), call detections follow a similar temporal pattern, such as the peaks in call rates occurring in July and December 2021 (**Figures 10, 11, 13**). Prior analyses from this area indicate Rice's whale calls can be detected up to 75 km on some occasions. Preliminary results from an acoustic tracking project, which leverages the deployment 1 data, finds that calls detected at site DCL are also detected at sites DCJ, DCK, DCM, DCN, DCO, and DCP, suggesting that at least part of the explanation for these shared temporal occurrence patterns is that they represent the same whales detected on multiple instruments. These patterns may also be indicative of broader population movement patterns.
- Preliminary results suggest potential seasonal movement patterns with higher detection rates of true long-moan calls at northern sites in spring and early summer and higher detection rates at southern sites in late summer to fall (**Figures 11, 13, 14, 15**). Spatial comparisons of percent of days per month present per site (**Figure 14**) and call detection rates per site (**Figure 15**) support the patterns observed in the time-series (**Figures 11, 13**). Additional data from the 3<sup>rd</sup> and 4<sup>th</sup> deployments may help determine whether these potential seasonal patterns are consistent beyond one year.
- Fewer long-moan calls were detected at the southern-most sites DCQ and DCR during the 2<sup>nd</sup> deployment than at the sites to the north of them (**Figures 11, 13**). This may indicate fewer whales occur here as the region approaches the known extents of the core Rice's whale habitat, or this may be the effect of masking due to higher levels of shipping and seismic airgun noise at these sites.

**Table 2.** Number of long-moan calls automatically detected and true calls validated per site during the three SoundTrap and HARP deployments at 18 sites over the May 2021 – Sept. 2022 period. No western Gulf long-moan variants were detected during this deployment. Shaded sites are from the offshore line. Numbers in italics from deployment 3 and site H deployment 2 indicate pre-validation results. Table will be updated when validations are complete.


Site	Deployment 1				Deployment 2				Deployment 3			
	Automated Detections	Validated Calls	Days Present (%)	Hours Present (%)	Automated Detections	Validated Calls	Days Present (%)	Hours Present (%)	Automated Detections	Validated Calls	Days Present (%)	Hours Present (%)
DCA	11,641	27	11 (7)	17 (0.5)	11,526	797	44 (31)	197 (5.8)	<i>14,389</i>	<i>n/a</i>	<i>159 (100)</i>	<i>2520 (67)</i>
DCB	16,927	10,208	80 (61)	1159 (36.7)	-	-	-	-	<i>15,354</i>	<i>n/a</i>	<i>151 (99)</i>	<i>2221 (61)</i>
DCC	-	-	-	-	4,654	3,239	88 (64)	855 (26.3)	<i>2,651</i>	<i>n/a</i>	<i>87 (100)</i>	<i>1219 (59)</i>
DCD	44,077	38,205	110 (96)	2321 (84.8)	14,905	11,978	91 (66)	1397 (42.5)	<i>35,856</i>	<i>n/a</i>	<i>150 (98)</i>	<i>2821 (77)</i>
DCE	5,455	1,107	87 (56)	461 (12.5)	1,509	0	0 (0)	0 (0.0)	<i>20,631</i>	<i>n/a</i>	<i>153 (101)</i>	<i>2660 (73)</i>
DCF	45,573	39,097	136 (99)	2858 (86.8)	23,345	16,407	121 (83)	1818 (52.2)	-	-	-	-
DCG	5,991	2,800	101 (89)	745 (27.5)	7,406	1,853	96 (71)	597 (18.5)	<i>1,045</i>	<i>n/a</i>	<i>107 (100)</i>	<i>678 (27)</i>
DCH	44,678	38,375	113 (76)	2364 (67.1)	<i>144,819</i>	<i>n/a</i>	<i>285 (100)</i>	<i>6608 (97)</i>	-	-	-	-
DCI	434	7	2 (11)	4 (1.0)	2,351	368	33 (29)	107 (3.9)	<i>26,219</i>	<i>n/a</i>	<i>164 (100)</i>	<i>3182 (82)</i>
DCJ	21,655	16,448	130 (95)	2074 (65.6)	50,735	35,186	134 (96)	2405 (72.5)	<i>57,943</i>	<i>n/a</i>	<i>171 (100)</i>	<i>3895 (96)</i>
DCK	14,452	8,124	120 (87)	1402 (42.5)	34,745	8,775	128 (91)	1519 (45.3)	<i>42,937</i>	<i>n/a</i>	<i>153 (100)</i>	<i>3277 (90)</i>
DCL	36,830	27,320	134 (93)	2278 (66.0)	73,353	53884	139 (95)	2747 (79.1)	-	-	-	-
DCM	26,372	10,459	135 (93)	1730 (50.1)	37,239	11,650	86 (83)	1513 (61.8)	<i>49,071</i>	<i>n/a</i>	<i>154 (100)</i>	<i>3390 (93)</i>
DCN	39,946	28,991	134 (94)	2355 (69.2)	69,766	48,068	136 (97)	2849 (84.9)	-	-	-	-
DCO	32,636	11,439	131 (90)	1763 (50.7)	-	-	-	-	-	-	-	-
DCP	19,321	13,946	96 (95)	1442 (60.0)	85,354	35,816	137 (94)	2541 (73.3)	<i>769</i>	<i>n/a</i>	<i>61 (100)</i>	<i>444 (31)</i>
DCQ	-	-	-	-	46,394	3,450	93 (89)	851 (33.8)	<i>5,596</i>	<i>n/a</i>	<i>125 (84)</i>	<i>516 (15)</i>
DCR	-	-	-	-	69,911	18,643	130 (95)	1842 (56.4)	-	-	-	-
<b>Total</b>	<b>365,988</b>	<b>246,553</b>			<b>678,012</b>	<b>212,651</b>			<b>272,461</b>	<b>0</b>		




**Figure 10.** Rice's whale long-moan call presence in 1-minute bins at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array over the May 2021 to September 2022 deployment period. Light blue marks represent verified false detections; dark blue marks represent true long-moan detections. Periods from April to September 2022 at all sites and the July 2021 to June 2022 period at site H, shown with only light blue marks, represent long-moan detections that have not yet been validated. Night time is indicated by gray hourglass shading. The darker gray blocked area represents periods without recording effort.




**Figure 11.** Daily total of Rice's whale long moan detections (light blue = autodetections, dark blue = verified long-moans, red line = false detections) at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array from May 2021 to September 2022. Gray blocks indicate periods without recording effort. Periods from April to September 2022 at all sites and the July 2021 to June 2022 period at site H, shown with only light blue bars, represent long-moan detections that have not yet been validated.




**Figure 12.** Example maximum detection ranges of long-moan calls as a function of ambient noise levels at 125 Hz for SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022. Estimated example detection ranges calculated assuming a call source level of 145 dB, detection threshold of 10 dB signal-to-noise, and geometric spreading transmission loss at  $17 \cdot \log R$ ; these assumptions require verification. Ambient noise data are currently undergoing QA/QC to salvage good quality recordings during some periods with questionable quality; poor quality recordings will be removed.



**Figure 13.** Weekly Rice's whale long-moan calls per hour of effort at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array from May 2021 to September 2022. Dark bars represent validated long-moan detections. Periods from April to September 2022 at all sites and the July 2021 to June 2022 period at site H have not yet been validated. Light gray blocks indicate periods without recording effort.



**Figure 14** Monthly Rice's whale long-moan presence, as percent of days, at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array over the May 2021 to September 2022 deployment period. Color represents percent of days with calls present. Open circles with no color represent no data; Site H from deployment 2, and all sites from deployment 3 are being validated and maps will be updated once complete. Coastline and 100m isobaths up to 500m are indicated.



**Figure 15.** Monthly average of Rice’s whale long-moan call detection rates at 18 sites in the De Soto Canyon sparse array over the May 2021 to September 2022 deployment period. Color represents call detections per hour. Open circles with no color represent no data; Site H from deployment 2, and all sites from deployment 3 are being validated and maps will be updated once complete. Coastline and 100m isobaths up to 500m are indicated.

## Rice's Whale Downsweep Sequence Calls

There were a total of 232,751 Rice's whale downsweep sequence automated detections in recordings from 18 sites in the DC array during the May 1, 2021 to September 21, 2022 period, ranging between 176 and 16,013 detections per deployment per SoundTrap site (**Table 3**). A total of 24,295 detections occurred on the 2<sup>nd</sup> HARP deployment at DCH, which spans both the 2<sup>nd</sup> and 3<sup>rd</sup> deployments of SoundTraps. Similar to previous analyses of the eight years of DC HARP recordings, downsweep detections were far less common than long-moan call detections (6x). Over the 1<sup>st</sup> and 2<sup>nd</sup> deployments at sites for which automated downsweep sequence detections have been manually validated, 26,071 of the 125,842 detections were validated as true downsweep sequence calls, with a range of 0 to 6,506 per deployment per site (**Table 3**). There were high false detection rates across sites over both deployments (51-100%) due both to the rarity of true calls at many sites and to confusion with ubiquitous seismic airgun pulses and a long-moan call-type that includes heavy amplitude modulation in the tail. Similar to long-moan calls, true downsweep sequence detections were higher at inshore sites than offshore sites across both deployments (**Table 3, Figures 16, 17**). Noise levels are higher at 100 Hz (the center of the downsweep sequence call range) than at 125 Hz (the center of the long-moan call range), and hence preliminary estimates of detection ranges based on ambient noise levels indicate detection ranges will be smaller for downsweep sequences (**Figure 18**) than for long-moans (**Figure 12**), assuming similar source levels among call types. Similar to long-moans, these detection ranges as a function of noise levels are generally higher at offshore sites than inshore sites, so this does not explain the decreased call detections at offshore sites (**Figure 17**). Unlike long-moan calls, true downsweep sequence detections were not found at all sites during either of the validated deployments, with several northern sites lacking detections in one or both deployments (**Table 3, Figures 16, 17, 19**). Similar to long-moan calls, there were suggestions of a seasonal movement between middle-habitat sites in the late spring to summer and southern sites in the late summer to fall (**Figures 17, 19, 20, 21**), though data from additional years (e.g. the 3<sup>rd</sup> and 4<sup>th</sup> deployments) are needed to verify if this is a consistent pattern across years.


- There were many fewer downsweep pulse sequence call detections (232,751) than long-moan detections (1,316,461) over the 18 passive acoustic sites (over 6x more long-moans), with an average of 4,195 downsweep sequence detections per site, and a range from 176 to 16,013 detections (**Tables 2, 3**).
- Validation of auto-detections of downsweep sequences has been completed on all sites from the 1<sup>st</sup> and 2<sup>nd</sup> deployments, except for the 2<sup>nd</sup> deployment HARP (DCH). True downsweep sequence detections did not occur at all sites during both validated deployments, with northern sites DCA, DCB, and DCE lacking detections during deployment 1 and sites DCA, DCC, DCD, DCE, and DCG lacking detections during deployment 2 (**Table 3, Figures 16, 17, 19**).
- Detections of true downsweep pulse sequences per deployment were higher at inshore sites (mean: 1,610; range: 0 to 6,506) than offshore sites (mean: 128; range: 0 to 810) (**Table 3**).
- Validation results from the 18 sites yielded false detection rates averaging 74% and 84% for the 1<sup>st</sup> and 2<sup>nd</sup> deployments, respectively, and ranging between 51% and 100% for the downsweep sequence detector per deployment across the 18 sites over the May 2021– April 2022 period.
- Highest false detection rates (90-100%) occurred at sites where downsweep calls were rarely or not at all detected, as well as at central, inshore sites (**Figure 17**) due to confusion with a long-moan call type that is amplitude modulated throughout. Confusion with ubiquitous airgun pulses also affected the detector.
- Preliminary results indicate that true Rice's whale downsweep sequences were present an average of 23% (range: 0 – 68%) of days per site, and were present in an average of 6.0% (range: 0 to 35%) of

recording hours per site during the two deployments from May 2021 to April 2022 deployment (**Table 2**). Percent of days and percent of hours present were generally higher at the inshore sites (average 35%, range 0-68% of days; average 11%, range 0-35% of hours) than the offshore sites (average 12%, range 0-48% of days; average 1.6%, range 0-8.4% of hours).


- Preliminary results of estimated maximum detection ranges of downsweep sequence calls as a function of ambient noise conditions at 100 Hz (assuming call source level of 145 dB, detection threshold of 10 dB SNR, and  $17\log R$  geometric spreading loss at all sites) show similar differences in detection ranges between offshore and inshore sites (generally larger at offshore sites where noise levels are lower; **Figure 18**), but these ranges are smaller for downsweep sequences than for long-moan calls as ambient noise levels are higher at 100 Hz than at 125 Hz (**Figure 12**). Future analyses will include sound propagation modeling using parabolic equations to better estimate site-specific transmission loss effects on detection ranges, and improved estimates of long-moan and downsweep call source levels based on acoustic localization and tracking.
- Preliminary results suggest potential seasonal movement patterns with higher detection rates of true long-moan calls at northern sites in spring and early summer and higher detection rates at southern sites in late summer to fall (**Figures 17, 19, 20, 21**). Spatial comparisons of percent of days per month present per site (**Figure 20**) and call detection rates per site (**Figure 21**) support the patterns observed in the time-series (**Figures 17, 19**). Additional data from the 3<sup>rd</sup> and 4<sup>th</sup> deployments may help determine whether these potential seasonal patterns are consistent beyond one year.

Table 3. Number of downsweep sequence calls automatically detected and true calls validated per site during the three SoundTrap and HARP deployments at 18 sites over the May 2021 – Sept. 2022 period. Numbers in italics from deployment 3 and site H deployment 2 indicate pre-validation results. Table will be updated when validations are complete.


Site	Deployment 1				Deployment 2				Deployment 3			
	Automated Detections	Validated Calls	Days Present (%)	Hours Present (%)	Automated Detections	Validated Calls	Days Present (%)	Hours Present (%)	Automated Detections	Validated Calls	Days Present (%)	Hours Present (%)
DCA	1,828	0	0 (0)	0 (0.0)	1,687	0	0 (0)	0 (0.0)	2,045	<i>n/a</i>	144 (91)	742 (20)
DCB	1,926	0	0 (0)	0 (0.0)	-	-	-	-	1,184	<i>n/a</i>	95 (62)	405 (11)
DCC	-	-	-	-	1,443	0	0 (0)	0 (0.0)	3,987	<i>n/a</i>	87 (100)	1411 (68)
DCD	8,610	57	3 (3)	14 (0.5)	3,073	0	0 (0)	0 (0.0)	18,465	<i>n/a</i>	142 (93)	2088 (57)
DCE	353	0	0 (0)	0 (0.0)	968	0	0 (0)	0 (0.0)	11,798	<i>n/a</i>	148 (97)	1769 (49)
DCF	7,166	1,068	30 (22)	209 (6.4)	6,645	418	19 (13)	94 (2.7)	-	-	-	-
DCG	176	20	3 (3)	6 (0.2)	2,149	0	0 (0)	0 (0.0)	1,269	<i>n/a</i>	100 (93)	535 (21)
DCH	16,013	6,506	101 (68)	1,233 (35.0)	24,295	<i>n/a</i>	283 (99)	3795 (56)	-	-	-	-
DCI	1,421	1	1 (6)	1 (0.2)	520	18	1 (1)	5 (0.2)	10,653	<i>n/a</i>	156 (95)	1400 (36)
DCJ	4,867	2,404	79 (58)	548 (17.3)	5,916	1,447	40 (29)	296 (8.9)	12,373	<i>n/a</i>	167 (98)	1797 (44)
DCK	2,628	810	66 (48)	276 (8.4)	4,227	149	30 (21)	73 (2.2)	10,638	<i>n/a</i>	145 (95)	1357 (37)
DCL	4,335	2,077	75 (52)	464 (13.5)	8,193	1,762	52 (36)	371 (11)	-	-	-	-
DCM	1,957	428	53 (37)	171 (4.9)	3,108	349	33 (32)	121 (4.9)	8,675	<i>n/a</i>	147 (95)	1332 (36)
DCN	4,422	1,319	63 (44)	361 (10.6)	7,052	2498	77 (55)	580 (17)	-	-	-	-
DCO	1,117	92	24 (17)	53 (1.5)	-	-	-	-	-	-	-	-
DCP	1,311	300	28 (28)	109 (4.5)	10,696	3,148	90 (62)	698 (20)	1,317	<i>n/a</i>	52 (85)	419 (29)
DCQ	-	-	-	-	4,328	54	19 (18)	30 (1.2)	210	<i>n/a</i>	26 (18)	93 (3)
DCR	-	-	-	-	7,707	1,146	67 (49)	312 (9.6)	-	-	-	-
<b>Total</b>	<b>58,130</b>	<b>15,082</b>			<b>92,007</b>	<b>10,989</b>			<b>82,614</b>	<b>-</b>		




**Figure 16.** Rice's whale downsweep sequence call presence in 1-minute bins at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array over the May 2021 to September 2022 deployment period. Light blue marks represent verified false detections; dark blue marks represent true long-moan detections. Periods from April to September 2022 at all sites and the July 2021 to June 2022 period at site H, shown with only light blue marks, represent long-moan detections that have not yet been validated.. The darker gray blocked area represents periods without recording effort.




**Figure 17.** Daily total of Rice's whale downsweep sequence detections (light blue = autodetections, dark blue = verified long-moans, red line = false detections) at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array from May 2021 to September 2022. Gray blocks indicate periods without recording effort. Periods from April to September 2022 at all sites and the July 2021 to June 2022 period at site H, shown with only light blue bars, represent long-moan detections that have not yet been validated.



**Figure 18.** Example maximum detection ranges of downsweep sequence calls as a function of ambient noise levels at 100 Hz for SoundTrap and HARP deployments at the De Soto Canyon (DC) array sites from May 2021 to September 2022. Estimated example detection ranges calculated assuming a call source level of 145 dB, detection threshold of 10 dB signal-to-noise, and geometric spreading transmission loss at  $17 \cdot \log R$ ; these assumptions require verification. Ambient noise data are currently undergoing QA/QC to salvage good quality recordings during some periods with questionable quality; poor quality recordings will be removed.



**Figure 19.** Weekly Rice's whale downsweep sequence calls per hour of effort at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array from May 2021 to September 2022. Dark bars represent validated long-moan detections. Periods from April to September 2022 at all sites and the July 2021 to June 2022 period at site H have not yet been validated. Light gray blocks indicate periods without recording effort.



**Figure 20.** Monthly Rice’s whale downsweep sequence presence, as percent of days, at 18 passive acoustic monitoring sites in the De Soto Canyon (DC) sparse array over the May 2021 to September 2022 deployment period. Color represents percent of days with calls present. Open circles with no color represent no data; Site H from deployment 2, and all sites from deployment 3 are being validated and maps will be updated once complete. Coastline and 100m isobaths up to 500m are indicated.




Figure 21. Monthly average of Rice's whale downsweep sequence call detection rates at 18 sites in the De Soto Canyon sparse array over the May 2021 to September 2022 deployment period. Color represents call detections per hour. Open circles with no color represent no data; Site H from deployment 2, and all sites from deployment 3 are being validated and maps will be updated once complete. Coastline and 100m isobaths up to 500m are indicated.

## **Vemco Acoustic Tag Detections**

Opportunistic data collection of transmissions from Vemco-acoustic-tagged fish and reptiles that pass by the moorings was possible due to the use of Vemco VR2AR acoustic releases. Tag detections occurred at all sites with a total of 1,090 transmissions received from 65 unique tagged individuals over the first three deployments. Vemco tag IDs have been uploaded to the iTag orphan tag database and shared with Innovasea, and 53 of the tagged individuals have been identified. The majority of individuals identified are great white sharks, while bull sharks, tiger sharks, scalloped hammerhead sharks, Atlantic tarpon, Crevall jack, greater amberjack, and leatherback turtles have also been identified.

## **Next Steps**

All data collection for this project is complete with the fourth and final deployment recovered in March 2023. The data processing phase of the Navy-funded deployments is over two-thirds complete with automated detections and validation steps complete for the first and second deployments. The automated detectors have been run on the third deployment recordings and the concurrently deployed HARP and are in the process of being validated. Analyses of the fourth deployment are planned for summer 2023 (funded by NOAA's OPR). Daily and monthly statistical distributions of sound pressure spectrum levels have been calculated for the first three deployments and will be calculated for the fourth deployment. Once all long-moan and downsweep sequence call detections have been validation for the third and fourth deployments, statistical analyses will be conducted to evaluate diel, seasonal, and spatial variation in call occurrence over the DC array, and to evaluate the impacts of varying ambient noise levels on call detection. These analyses will provide crucially important data for understanding how Rice's whales are utilizing the core habitat throughout the course of the year and whether they exhibited seasonal movement patterns throughout the year. Understanding seasonal and internannual distribution of Rice's whales throughout the core habitat will improve understanding of potential impact of human activities on these whales and assist in developing effective mitigation measures as needed.

## **Data Leveraging and Future Opportunities**

Several analytic efforts are underway that are leveraging data collected under this project to further improve our understanding of Rice's whale seasonal and interannual density and additional analytical projects could be developed to improve our understanding of the oceanographic and anthropogenic factors driving these patterns. To convert occurrence and distribution results into animal density, more information is needed on calling frequency and detection ranges and how they vary over time and space due to sound propagation conditions and ambient noise levels. Expanded analytical methods are being developed to leverage the SoundTrap array data to fill this need. While the sparse array was designed to ensure near-complete acoustic coverage of the Rice's whale core habitat, assuming minimum detection distances of 20 km based on prior findings, to understand seasonal distribution and movement patterns, further evaluation found calls are commonly detected on multiple neighboring instruments in the array. Under funding from NOAA's Ocean Acoustics Program, whale calls are being localized and tracked using time-difference-of-arrival (TDOA) methods, to obtain information on call detection distances, source levels, call rates, and swimming behavior, with automated localization algorithms in development. Given the occurrence of the same call on multiple instruments, spatially-explicit capture-recapture (SECR) analyses could be conducted to estimate density and detection distances, providing complementary results to evaluate detection distances and how they change over time along with density estimates. Further, to permit and mitigate impacts from anthropogenic activities in this core habitat area where these endangered whales consistently occur year-round, predictive habitat models describing the factors driving spatio-temporal occurrence will be important to assess and predict when and where the whales might be found to determine if we can better predict finer-

scale spatial occurrence. As detection distances, and how they vary over time, are better understood with the ongoing tracking analyses, future studies could combine acoustic detections with oceanographic data to evaluate if variation in the position of the Loop Current and its eddies, or Mississippi River outflow, may impact animal distribution as well as sound propagation conditions and associated call detection distances. Developing predictive habitat models incorporating environmental proxies of prey occurrence, ambient noise levels, and modeled detection distances with passive acoustic detections as the response variable will help determine which dynamic factors drive the occurrence of calling Rice's whales throughout the core habitat, as needed to mitigate potential impacts from anthropogenic activities occurring in the area.

Finally, the data collection component of this project covers nearly two years to evaluate seasonal changes in Rice's whale distribution throughout the core habitat. The anticipated results will characterize spatio-temporal variation for the May 2021 to March 2023 period. To make further inferences about whether these trends represent general seasonal changes, we suggest a minimum of three years of data collection to evaluate consistency in seasonal cycles over time. At this time it remains unknown whether factors driving temporal variation in Rice's whale occurrence and distribution follow typical four-season cycles or are more nuanced with respect to oceanographic conditions including the position of the Loop Current and its eddies and variation in Mississippi River outflow. Three years of broad coverage passive acoustic data collection would provide the information needed to assess generality of the 2021-2023 results and would yield a more robust dataset for developing the predictive habitat models described above.

## References

- Garrison, L. P., Ortega-Ortiz, J., and Rappucci, G. (2020). "Abundance of Marine Mammals in Waters of the U.S. Gulf of Mexico during the Summers of 2017 and 2018," in Southeast Fisheries Science Center Reference Document PRBD-2020-07 (Miami, FL), p. 56.
- Hildebrand JA (2009) Anthropogenic and natural sources of ambient noise in the ocean. *Marine Ecology Progress Series* 395:5-20
- Mellinger DK, Clark CW (2000) Recognizing transient low-frequency whale sounds by spectrogram correlation. *Journal of the Acoustical Society of America* 107:12
- Rice AN, Palmer KJ, Tielens JT, Muirhead CA, Clark CW (2014) Potential Bryde's whale (*Balaenoptera edeni*) calls recorded in the northern Gulf of Mexico. *The Journal of the Acoustical Society of America* 135:3066-3076
- Rosel, P. E., Wilcox, L. A., Yamada, T. K., and Mullin, K. D. (2021). "A new species of baleen whale (*Balaenoptera*) from the Gulf of Mexico, with a review of its geographic distribution," *Marine Mammal Science*. 10.1111/mms.12776
- Rosel, P.E. and Garrison, L.P. (2022). Rice's whale core distribution map Version 7 June 2019. NMFS Southeast Fisheries Science Center Reference Document MMTD-2022-01.
- Širović A, Bassett HR, Johnson SC, Wiggins SM, Hildebrand JA (2014) Bryde's whale calls recorded in the Gulf of Mexico. *Marine Mammal Science* 30:399-409
- Soldevilla, M.S., Debich, A.J., Garrison, L.P., Hildebrand, J.A., & Wiggins, S.M. (2022a). Rice's whales in the northwestern Gulf of Mexico: Call variation and occurrence beyond the known core habitat. *Endangered Species Research*, 48:155-174
- Soldevilla MS, Hildebrand JA, Frasier KE, Aichinger Dias L and others (2017) Spatial Distribution and Dive Behavior of Gulf of Mexico Bryde's Whales: Potential Risk for Vessel Strikes and Fisheries Interactions. *Endangered Species Research* 32:533-550
- Soldevilla, M.S., Ternus, K., Cook, A., Hildebrand, J.A., Frasier, K.E., Martinez, A., & Garrison, L.P. (2022b). Acoustic localization, validation, and characterization of Rice's whale calls. *Journal of the Acoustical Society of America*. 151:4264-4278
- Wiggins SM, Hall J, Thayre BJ, Hildebrand JA (2016) Gulf of Mexico low-frequency ocean soundscape dominated by airguns. *Journal of the Acoustical Society of America* 140:176-183
- Wiggins SM, Hildebrand JA (2007) High-frequency Acoustic Recording Package (HARP) for broadband, long-term marine mammal monitoring. *International Symposium on Underwater Technology 2007 and International Workshop on Scientific Use of Submarine Cables and Related Technologies* 2007:551-557